



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

DECLARATION

I, Dr. R.H. Walter, BSc., PhD., EurChem., CChem., MRSC., MITI., translator to Taylor & Meyer of 20 Kingsmead Road, London SW2 3JD, England, do solemnly and sincerely declare as follows:

1. That I am well acquainted with the English and German languages;
2. That the following is a true translation made by me into the English language of International Patent Application No. PCT/EP02/08729;
3. That all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true;

and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardise the validity of the application or any patent issued thereon.

Signed, this *13<sup>th</sup>* day of *January* 2006,

*R.H. Walter*

High Wycombe, Bucks., HP10 9RH, England

Patent application in the name of Grünenthal GmbH, D-52078  
Aachen  
(in-house reference G 3068)

5

Salts of substituted 1,2,3,4-tetrahydroquinoline-2-  
carboxylic acid derivatives

10 The present invention relates to salts of substituted  
1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivatives,  
to processes for their preparation, to drugs containing  
these compounds and to their use for the preparation of  
drugs for specific indications, especially for the  
treatment of pain.

15

The treatment of chronic and non-chronic pain conditions is  
of great importance in medicine. There is a worldwide need  
for highly effective pain therapies for the patient-  
orientated and targeted treatment of chronic and non-  
20 chronic pain conditions, this being understood as meaning  
the successful and satisfactory treatment of pain for the  
patient. This manifests itself in the large number of  
scientific studies which have recently appeared in the  
field of applied analgesis or fundamental research into  
25 nociception.

Conventional opioids such as morphine are highly effective  
in the therapy of intense to very intense pain. However,  
their use is limited by the known side effects, e.g.  
30 respiratory depression, vomiting, sedation, constipation  
and the development of tolerance. Also, they are less  
effective in cases of neuropathic or incidental pain, as  
suffered by tumour patients in particular.

35 Opioids develop their analgesic effect by binding to  
membrane-based receptors belonging to the family of the

so-called G protein coupled receptors. The biochemical and pharmacological characterization of subtypes of these receptors has now aroused the hope that subtype-specific opioids possess a different action/side effect profile from  
5 that of e.g. morphine. Further pharmacological studies have since suggested the probable existence of several subtypes of these opioid receptors ( $\mu_1$ ,  $\mu_2$ ,  $\kappa_1$ ,  $\kappa_2$ ,  $\kappa_3$ ,  $\delta_1$  and  $\delta_2$ ).

10 In addition, there are other receptors and ion channels which play a substantial part in the system of the development and transmission of pain. The NMDA ion channel is of particular importance here because a substantial  
15 proportion of the communication of synapses passes through it. The calcium ion exchange between a neuronal cell and its environment is controlled by this channel.

Knowledge about the physiological significance of ion channel-selective substances has been gained by the  
20 development of the patch-clamp technique. The action of NMDA antagonists on the influx of calcium ions into the interior of the cell can be unambiguously demonstrated in this way. It has also been shown that these substances possess an independent antinociceptive potential (e.g.  
25 ketamine). It is important here that the mechanism of action is quite different than e.g. in the case of opiates, because NMDA antagonists intervene directly in the decisive calcium balance of the cells during the transmission of pain. This affords the possibility for the first time of  
30 treating neuropathic forms of pain successfully.


Various NMDA antagonists, in this case involving tetrahydroquinoline derivatives, have already been described in the articles J. Med. Chem. (1992) 35, 1954-  
35 1968, J. Med. Chem. (1992) 35, 1942-1953 and Med. Chem. Res. (1991) 1, 64-73, and in patent applications EP 386

839, WO 97/12879 A1, WO 98/07704 A1 and WO 98/42673 A1.

Said references, especially the patent applications, gave a large number of possible indications, including pain therapy. However, the efficacy and applicability of these substances is still open to improvement, so there is a need for other substances here.

One object of the invention was to provide analgesically effective substances, especially NMDA antagonists, which are suitable for pain therapy, including chronic and neuropathic pain in particular. Furthermore, these substances should exhibit a minimum of side effects, e.g. nausea, vomiting, dependence, respiratory depression or constipation.

Accordingly, the invention provides 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivatives of general formula I in the form shown, or in the form of their acids or their bases, or in the form of their salts, especially the physiologically acceptable salts, or in the form of their solvates, especially the hydrates, particularly in the form of their physiologically acceptable salts with cations or bases or with anions or acids, optionally in the form of their racemates or their pure stereoisomers, especially enantiomers or diastereoisomers, or in the form of mixtures of the stereoisomers, especially the enantiomers or diastereoisomers, in any desired mixing ratio:




I,

in which

**either**

- 5 **R<sup>1</sup>** and **R<sup>2</sup>** together form the following, each of which is monosubstituted or polysubstituted or unsubstituted:

- (CH<sub>2</sub>)<sub>n</sub>-, where n = 3-10,  
 -CH=CH-CH<sub>2</sub>-, -CH<sub>2</sub>-CH=CH-,  
 10 -CH=CH-CH<sub>2</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-CH<sub>2</sub>-CH=CH-,  
 -CH<sub>2</sub>-CH=CH-CH<sub>2</sub>-,  
 -CH<sub>2</sub>-CH=CH-CH<sub>2</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-CH<sub>2</sub>-CH=CH-CH<sub>2</sub>-,  
 -CH<sub>2</sub>-CH<sub>2</sub>-CH=CH-CH<sub>2</sub>-CH<sub>2</sub>-,  
 -O-CH<sub>2</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-CH<sub>2</sub>-O-,  
 15 -O-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-O-,  
 -CH<sub>2</sub>-O-CH<sub>2</sub>-,  
 -CH<sub>2</sub>-CH<sub>2</sub>-O-CH<sub>2</sub>-, -CH<sub>2</sub>-O-CH<sub>2</sub>-CH<sub>2</sub>-,



$R^3$  is selected from

- 5 H;  $C_1$ - $C_{18}$ -alkyl,  $C_2$ - $C_{18}$ -alkenyl or  $C_2$ - $C_{18}$ -alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted;  $C_3$ - $C_8$ -cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted, or a
- 10 corresponding heterocycle in which at least one ring C atom is replaced by N, S or O; alkylaryl or alkylheteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted; and aryl or heteroaryl, each of which is monosubstituted or
- 15 polysubstituted or unsubstituted;

naphthyl or 2-naphthyl, 2-, 3- or 4-pyridyl or thiazolyl;

SR<sup>10</sup>, where R<sup>10</sup> is selected from

5

aryl or heteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted;

10

C(O)NR<sup>11</sup>R<sup>12</sup>, C(O)NR<sup>11</sup>NR<sup>12</sup>R<sup>13</sup>, C(NR<sup>11</sup>)NR<sup>12</sup>R<sup>13</sup>, C(S)NR<sup>11</sup>R<sup>12</sup> or C(S)NR<sup>11</sup>NR<sup>12</sup>R<sup>13</sup>, where R<sup>11</sup>, R<sup>12</sup> and R<sup>13</sup> independently of one another are selected from

15

H; C<sub>1</sub>-C<sub>18</sub>-alkyl, C<sub>2</sub>-C<sub>18</sub>-alkenyl or C<sub>2</sub>-C<sub>18</sub>-alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted; C<sub>3</sub>-C<sub>8</sub>-cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted, or a corresponding heterocycle in which at least one ring C atom is replaced by S, O or N; alkylaryl or alkylheteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted; and aryl or heteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted;

20

25

30

R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> independently of one another are selected from

35

H; F; Cl; Br; I; CN; NO<sub>2</sub>; and C<sub>1</sub>-C<sub>10</sub>-alkyl, C<sub>2</sub>-C<sub>10</sub>-alkenyl or C<sub>2</sub>-C<sub>10</sub>-alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted;

$R^4$  is selected from

$R^{4a}$  or  $ZR^{4a}$ , where  $Z = C_1-C_6$ -alkyl,  $C_2-C_6$ -alkenyl or  
5  $C_2-C_6$ -alkynyl, each of which is branched or unbranched  
and monosubstituted or polysubstituted or  
unsubstituted; and  $R^{4a}$  is selected from

10 H;  $C_1-C_{12}$ -alkyl,  $C_2-C_{12}$ -alkenyl or  $C_2-C_{12}$ -alkynyl,  
each of which is branched or unbranched and  
monosubstituted or polysubstituted or  
unsubstituted;  $C_3-C_8$ -cycloalkyl which is saturated  
or unsaturated and monosubstituted or  
polysubstituted or unsubstituted, or a  
15 corresponding heterocycle in which at least one  
ring C atom is replaced by S, O or N; and aryl or  
heteroaryl, each of which is monosubstituted or  
polysubstituted or unsubstituted;

20  $C(O)R^9$ ,  $C(O)OR^9$ ,  $C(S)R^9$ ,  $C(S)OR^9$  or  $S(O_2)R^9$ , where  
 $R^9$  is selected from

25 H;  $C_1-C_{10}$ -alkyl,  $C_2-C_{10}$ -alkenyl or  
 $C_2-C_{10}$ -alkynyl, each of which is branched or  
unbranched and monosubstituted or  
polysubstituted or unsubstituted;  $C_3-C_8$ -  
cycloalkyl which is saturated or unsaturated  
and monosubstituted or polysubstituted or  
unsubstituted, or a corresponding  
30 heterocycle in which at least one ring C  
atom is replaced by S, O or N; alkylaryl or  
alkylheteroaryl, each of which is  
monosubstituted or polysubstituted or  
unsubstituted; and aryl or heteroaryl, each  
of which is monosubstituted or  
35 polysubstituted or unsubstituted, especially  
phenethyl, 1-adamantyl, 2-adamantyl, 1-



OR<sup>14</sup>, OC(O)R<sup>14</sup>, OC(S)R<sup>14</sup>, C(O)R<sup>14</sup>, C(O)OR<sup>14</sup>, C(S)R<sup>14</sup>, C(S)OR<sup>14</sup>, SR<sup>14</sup>, S(O)R<sup>14</sup> or S(O<sub>2</sub>)R<sup>14</sup>, where R<sup>14</sup> is selected from

5 H; C<sub>1</sub>-C<sub>10</sub>-alkyl, C<sub>2</sub>-C<sub>10</sub>-alkenyl or C<sub>2</sub>-C<sub>10</sub>-alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted; C<sub>3</sub>-C<sub>8</sub>-cycloalkyl which is saturated or unsaturated and monosubstituted or  
10 polysubstituted or unsubstituted, or a corresponding heterocycle in which at least one ring C atom is replaced by S, O or N; alkylaryl or alkylheteroaryl, each of which is monosubstituted or polysubstituted or  
15 unsubstituted; and aryl or heteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted;

NR<sup>15</sup>R<sup>16</sup>, NR<sup>15</sup>C(O)R<sup>16</sup>, C(NR<sup>15</sup>)NR<sup>16</sup>R<sup>17</sup>, NR<sup>15</sup>C(S)R<sup>16</sup>,  
20 C(S)NR<sup>15</sup>R<sup>16</sup>, C(S)NR<sup>15</sup>NR<sup>16</sup>R<sup>17</sup> or S(O<sub>2</sub>)NR<sup>15</sup>R<sup>16</sup>, where R<sup>15</sup>, R<sup>16</sup> and R<sup>17</sup> independently of one another are selected from

H; O; C<sub>1</sub>-C<sub>18</sub>-alkyl, C<sub>2</sub>-C<sub>18</sub>-alkenyl or C<sub>2</sub>-C<sub>18</sub>-alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or  
25 unsubstituted; C<sub>3</sub>-C<sub>8</sub>-cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted, or a corresponding heterocycle in which at least one ring C atom is replaced by S, O or N; alkylaryl or alkylheteroaryl, each of which is monosubstituted or polysubstituted or  
30 unsubstituted; and aryl or heteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted;  
35

or

5  $R^{15}$  and  $R^{16}$  or  $R^{16}$  and  $R^{17}$  together form a  $C_3$ - $C_8$ -cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted, or a corresponding heterocycle in which at least one ring C atom is replaced by S, O or N; or


10  $R^5$  and  $R^6$ ,  $R^6$  and  $R^7$  or  $R^7$  and  $R^8$  together form

$=CR^{18}-CH=CH-CH=$  or  $=CH-CR^{18}=CH-CH=$ , where  $R^{18}$  is selected from

15 H; F; Cl; Br; I; OH; and  $C_1$ - $C_{10}$ -alkyl,  $C_2$ - $C_{10}$ -alkenyl or  $C_2$ - $C_{10}$ -alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted,

20 with the proviso that

if  $R^1$  and  $R^2$  together form  $-CH=CH-CH_2-$  or



25


and  $R^3$  is (-)-p-menthan-3-ol, especially menthol or borneol,  $R^7 \neq Cl$  and  $R^5$ ,  $R^6$  and  $R^8 \neq H$  simultaneously,

30 if  $R^1$  and  $R^2$  together form  $-CH=CH-CH_2-$  and  $R^3$  is  $CH_3$ ,  $R^7 \neq H$ , Cl or  $OCH_3$  and  $R^5$ ,  $R^6$  and  $R^8 \neq H$  simultaneously,

if  $R^{1b}$  and  $R^{2a}$  together form  $-CH=CH-CH_2-$  and  $R^3$  is H,  $R^7 \neq OCH_3$  or  $C(O)NH_2$  and  $R^5$ ,  $R^6$  and  $R^8 \neq H$ ,  $R^5$  and  $R^7 \neq CH_3$  and  $R^6$  and  $R^8 \neq H$ , or  $R^5 \neq OCH_3$  and  $R^6$ ,  $R^7$  and  $R^8 \neq H$

simultaneously, or

if  $R^{1b}$  and  $R^{2a}$  together form



or  $-O-CH_2-CH_2-$  and  $R^3$  is  $C_2H_5$ ,  $R^7 \neq H$ ,  $Cl$ ,  $CH_3$ ,  $OCH_3$  or  $NO_2$  and  $R^5$ ,  $R^6$  and  $R^8 \neq H$ , or  $R^5 \neq NO_2$  and  $R^6$ ,  $R^7$  and  $R^8 \neq H$  simultaneously;

**or**

$R^1$  is selected from

$C_1-C_{10}$ -alkyl which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted;  $C_3-C_8$ -cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted, or a corresponding heterocycle in which at least one ring C atom is replaced by S, O or N; alkylaryl which is monosubstituted or polysubstituted or unsubstituted; and aryl which is monosubstituted or polysubstituted or unsubstituted;

$OR^{19}$ ,  $SR^{19}$  or  $SO_2R^{19}$ , where  $R^{19}$  is selected from

$C_1-C_{10}$ -alkyl,  $C_2-C_{10}$ -alkenyl or  $C_2-C_{10}$ -alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted;  $C_3-C_8$ -cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted, or a corresponding heterocycle in which at least one ring C atom is replaced by S, O or N; and

alkylaryl, aryl, alkylheteroaryl or heteroaryl,  
each of which is monosubstituted or  
polysubstituted or unsubstituted;

5  $R^2$  is selected from

H;  $C_1$ - $C_{10}$ -alkyl which is branched or unbranched and  
monosubstituted or polysubstituted or unsubstituted;  
and phenyl which is monosubstituted or polysubstituted  
10 or unsubstituted, where if  $R^2$  is phenyl,  $R^1$  must be  
aryl, O-aryl or S-aryl;

$R^3$  is selected from

15 H;  $C_1$ - $C_{18}$ -alkyl,  $C_2$ - $C_{18}$ -alkenyl or  $C_2$ - $C_{18}$ -alkynyl, each  
of which is branched or unbranched and monosubstituted  
or polysubstituted or unsubstituted;  $C_3$ - $C_8$ -cycloalkyl  
which is saturated or unsaturated and monosubstituted  
or polysubstituted or unsubstituted, or a  
20 corresponding heterocycle in which at least one ring C  
atom is replaced by N, S or O; alkylaryl or  
alkylheteroaryl, each of which is monosubstituted or  
polysubstituted or unsubstituted; and aryl or  
heteroaryl, each of which is monosubstituted or  
25 polysubstituted or unsubstituted;

$R^4$  is selected from

30  $R^{4a}$  or  $ZR^{4a}$ , where  $Z = C_1$ - $C_6$ -alkyl,  $C_2$ - $C_6$ -alkenyl or  
 $C_2$ - $C_6$ -alkynyl, each of which is branched or unbranched  
and monosubstituted or polysubstituted or  
unsubstituted; and  $R^{4a}$  is selected from

35 H;  $C_1$ - $C_{12}$ -alkyl,  $C_2$ - $C_{12}$ -alkenyl or  $C_2$ - $C_{12}$ -alkynyl,  
each of which is branched or unbranched and  
monosubstituted or polysubstituted or

unsubstituted; C<sub>3</sub>-C<sub>8</sub>-cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted, or a corresponding heterocycle in which at least one ring C atom is replaced by S, O or N; and aryl or heteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted;

C(O)R<sup>9</sup>, C(O)OR<sup>9</sup>, C(S)R<sup>9</sup>, C(S)OR<sup>9</sup> or S(O<sub>2</sub>)R<sup>9</sup>, where R<sup>9</sup> is selected from

H; C<sub>1</sub>-C<sub>10</sub>-alkyl, C<sub>2</sub>-C<sub>10</sub>-alkenyl or C<sub>2</sub>-C<sub>10</sub>-alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted; C<sub>3</sub>-C<sub>8</sub>-cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted, or a corresponding heterocycle in which at least one ring C atom is replaced by S, O or N; alkylaryl or alkylheteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted; and aryl or heteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted, especially phenethyl, 1-adamantyl, 2-adamantyl, 1-naphthyl or 2-naphthyl, 2-, 3- or 4-pyridyl or thiazolyl;

SR<sup>10</sup>, where R<sup>10</sup> is selected from

aryl or heteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted;

C(O)NR<sup>11</sup>R<sup>12</sup>, C(O)NR<sup>11</sup>NR<sup>12</sup>R<sup>13</sup>, C(NR<sup>11</sup>)NR<sup>12</sup>R<sup>13</sup>,

$C(S)NR^{11}R^{12}$  or  $C(S)NR^{11}NR^{12}R^{13}$ , where  $R^{11}$ ,  $R^{12}$  and  $R^{13}$  independently of one another are selected from

5 H;  $C_1$ - $C_{18}$ -alkyl,  $C_2$ - $C_{18}$ -alkenyl or  $C_2$ - $C_{18}$ -alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted;  $C_3$ - $C_8$ -cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or  
10 unsubstituted, or a corresponding heterocycle in which at least one ring C atom is replaced by S, O or N; alkylaryl or alkylheteroaryl, each of which is monosubstituted or polysubstituted or  
15 unsubstituted; and aryl or heteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted;

20  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  independently of one another are selected from

H; F; Cl; Br; I; CN;  $NO_2$ ; and  $C_1$ - $C_{10}$ -alkyl,  $C_2$ - $C_{10}$ -alkenyl or  $C_2$ - $C_{10}$ -alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or  
25 unsubstituted;

$OR^{14}$ ,  $OC(O)R^{14}$ ,  $OC(S)R^{14}$ ,  $C(O)R^{14}$ ,  $C(O)OR^{14}$ ,  $C(S)R^{14}$ ,  $C(S)OR^{14}$ ,  $SR^{14}$ ,  $S(O)R^{14}$  or  $S(O_2)R^{14}$ , where  $R^{14}$  is selected from

30 H;  $C_1$ - $C_{10}$ -alkyl,  $C_2$ - $C_{10}$ -alkenyl or  $C_2$ - $C_{10}$ -alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted;  $C_3$ - $C_8$ -cycloalkyl which is saturated or unsaturated and monosubstituted or  
35 polysubstituted or unsubstituted, or a

corresponding heterocycle in which at least one ring C atom is replaced by S, O or N; alkylaryl or alkylheteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted; and aryl or heteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted;

$\text{NR}^{15}\text{R}^{16}$ ,  $\text{NR}^{15}\text{C}(\text{O})\text{R}^{16}$ ,  $\text{C}(\text{NR}^{15})\text{NR}^{16}\text{R}^{17}$ ,  $\text{NR}^{15}\text{C}(\text{S})\text{R}^{16}$ ,  $\text{C}(\text{S})\text{NR}^{15}\text{R}^{16}$ ,  $\text{C}(\text{S})\text{NR}^{15}\text{NR}^{16}\text{R}^{17}$  or  $\text{S}(\text{O}_2)\text{NR}^{15}\text{R}^{16}$ , where  $\text{R}^{15}$ ,  $\text{R}^{16}$  and  $\text{R}^{17}$  independently of one another are selected from

H; O;  $\text{C}_1$ - $\text{C}_{18}$ -alkyl,  $\text{C}_2$ - $\text{C}_{18}$ -alkenyl or  $\text{C}_2$ - $\text{C}_{18}$ -alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted;  $\text{C}_3$ - $\text{C}_8$ -cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted, or a corresponding heterocycle in which at least one ring C atom is replaced by S, O or N; alkylaryl or alkylheteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted; and aryl or heteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted;

or

$\text{R}^{15}$  and  $\text{R}^{16}$  or  $\text{R}^{16}$  and  $\text{R}^{17}$  together form a  $\text{C}_3$ - $\text{C}_8$ -cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted, or a corresponding heterocycle in which at least one C atom is replaced by S, O or N; or

$\text{R}^5$  and  $\text{R}^6$ ,  $\text{R}^6$  and  $\text{R}^7$  or  $\text{R}^7$  and  $\text{R}^8$  together form

$=\text{CR}^{18}-\text{CH}=\text{CH}-\text{CH}=\text{}$  or  $=\text{CH}-\text{CR}^{18}=\text{CH}-\text{CH}=\text{}$ , where  $\text{R}^{18}$  is selected from

5 H; F; Cl; Br; I; OH; and  $\text{C}_1$ - $\text{C}_{10}$ -alkyl,  $\text{C}_2$ - $\text{C}_{10}$ -alkenyl or  $\text{C}_2$ - $\text{C}_{10}$ -alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted,

with the proviso that

10

if  $\text{R}^4$ ,  $\text{R}^6$ ,  $\text{R}^7$  and  $\text{R}^8 = \text{H}$ ,

- $\text{R}^1 \neq \text{CH}_3$ ,  $\text{R}^3 \neq \text{H}$  or  $\text{CH}_3$  and  $\text{R}^2$  and  $\text{R}^5 \neq \text{H}$  simultaneously;  
or

15 •  $\text{R}^1 \neq$  unsubstituted phenyl,  $\text{R}^3 \neq \text{C}_2\text{H}_5$  and  $\text{R}^2$  and  $\text{R}^5 \neq \text{H}$  simultaneously;

if  $\text{R}^4$ ,  $\text{R}^5$ ,  $\text{R}^6$  and  $\text{R}^8 = \text{H}$ ,

- $\text{R}^1 \neq$  S-phenyl,  $\text{R}^2 \neq \text{H}$ ,  $\text{R}^7 \neq \text{Cl}$  and  $\text{R}^3 \neq \text{CH}_3$  simultaneously;  
or

20 •  $\text{R}^1 \neq$  S-2-pyridinyl,  $\text{R}^2 \neq \text{CH}_3$ ,  $\text{R}^7 \neq \text{OCH}_3$  and  $\text{R}^3 \neq -\text{CH}_3-\text{CH}=\text{CH}_2$  simultaneously; or

if  $\text{R}^2$ ,  $\text{R}^4$ ,  $\text{R}^5$  and  $\text{R}^7 = \text{H}$  and  $\text{R}^6$  and  $\text{R}^8 = \text{Cl}$ ,

- $\text{R}^1 \neq$  dioxalane and  $\text{R}^3 \neq -\text{CH}_2-\text{CH}_2-\text{OH}$  simultaneously.

25

The 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivatives or their salts according to the invention exhibit a pronounced analgesic action and are also NMDA antagonists which selectively attack at the glycine binding site, and.

30

In terms of the present invention, alkyl or cycloalkyl radicals are understood as meaning saturated and unsaturated (but not aromatic), branched, unbranched and cyclic hydrocarbons which can be unsubstituted or monosubstituted or polysubstituted.  $\text{C}_{1-2}$ -alkyl is Cl- or

35



C2-alkyl, C<sub>1-3</sub>-alkyl is C1-, C2- or C3-alkyl, C<sub>1-4</sub>-alkyl is C1-, C2-, C3- or C4-alkyl, C<sub>1-5</sub>-alkyl is C1-, C2-, C3-, C4- or C5-alkyl, C<sub>1-6</sub>-alkyl is C1-, C2-, C3-, C4-, C5- or C6-alkyl, C<sub>1-7</sub>-alkyl is C1-, C2-, C3-, C4-, C5-, C6- or C7-alkyl, C<sub>1-8</sub>-alkyl is C1-, C2-, C3-, C4-, C5-, C6-, C7- or C8-alkyl, C<sub>1-10</sub>-alkyl is C1-, C2-, C3-, C4-, C5-, C6-, C7-, C8-, C9- or C10-alkyl and C<sub>1-18</sub>-alkyl is C1-, C2-, C3-, C4-, C5-, C6-, C7-, C8-, C9-, C10-, C11-, C12-, C13-, C14-, C15-, C16-, C17- or C18-alkyl. Also, C<sub>3-4</sub>-cycloalkyl is C3- or C4-cycloalkyl, C<sub>3-5</sub>-cycloalkyl is C3-, C4- or C5-cycloalkyl, C<sub>3-6</sub>-cycloalkyl is C3-, C4-, C5- or C6-cycloalkyl, C<sub>3-7</sub>-cycloalkyl is C3-, C4-, C5-, C6- or C7-cycloalkyl, C<sub>3-8</sub>-cycloalkyl is C3-, C4-, C5-, C6-, C7- or C8-cycloalkyl, C<sub>4-5</sub>-cycloalkyl is C4- or C5-cycloalkyl, C<sub>4-6</sub>-cycloalkyl is C4-, C5- or C6-cycloalkyl, C<sub>4-7</sub>-cycloalkyl is C4-, C5-, C6- or C7-cycloalkyl, C<sub>5-6</sub>-cycloalkyl is C5- or C6-cycloalkyl and C<sub>5-7</sub>-cycloalkyl is C5-, C6- or C7-cycloalkyl. 'Cycloalkyl' also embraces saturated cycloalkyls in which one or 2 carbon atoms are replaced by the heteroatom S, N or O. However, 'cycloalkyl' also includes especially monounsaturated or polyunsaturated and preferably monounsaturated cycloalkyls without a heteroatom in the ring as long as the cycloalkyl is not an aromatic system. The alkyl or cycloalkyl radicals are preferably methyl, ethyl, vinyl (ethenyl), propyl, allyl (2-propenyl), 1-propynyl, methylethyl, butyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl, pentyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 2,2-dimethylpropyl, hexyl, 1-methylpentyl, cyclopropyl, 2-methylcyclopropyl, cyclopropylmethyl, cyclobutyl, cyclopentyl, cyclopentylmethyl, cyclohexyl, cycloheptyl or cyclooctyl, but also adamantyl, CHF<sub>2</sub>, CF<sub>3</sub> or CH<sub>2</sub>OH, as well as pyrazolinone, oxopyrazolinone, 1,4-dioxane or dioxolane.

In the context of alkyl and cycloalkyl, 'substituted' in terms of the present invention is understood as meaning the

substitution of a hydrogen radical by F, Cl, Br, I, NH<sub>2</sub>, SH or OH, 'polysubstituted' radicals being understood as meaning that the substitution takes place both on different atoms and on the same atoms several times with the same or  
5 different substituents, for example three times on the same C atom, as in the case of CF<sub>3</sub>, or at different sites, as in the case of -CH(OH)-CH=CH-CHCl<sub>2</sub>. Particularly preferred substituents here are F, Cl and OH.

- 10 '(CH<sub>2</sub>)<sub>3-6</sub>' is understood as meaning -CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-,  
-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>- and  
-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-, '(CH<sub>2</sub>)<sub>1-4</sub>' is understood as meaning  
-CH<sub>2</sub>-, -CH<sub>2</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>- and -CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-, etc.
- 15 Aryl radicals are understood as meaning ring systems having at least one aromatic ring, but without heteroatoms in only one of the rings. Examples are phenyl, naphthyl, fluoroanthenyl, fluorenyl, tetralinyl or indanyl radicals, especially 9H-fluorenyl or anthracenyl radicals, which can  
20 be unsubstituted or monosubstituted or polysubstituted.

Heteroaryl radicals are understood as meaning heterocyclic ring systems having at least one unsaturated ring which contain one or more heteroatoms from the group comprising  
25 nitrogen, oxygen and/or sulfur, and which can also be monosubstituted or polysubstituted. Examples of the heteroaryl group which may be mentioned are furan, benzofuran, thiophene, benzothiophene, pyrrole, pyridine, pyrimidine, pyrazine, quinoline, isoquinoline, phthalazine,  
30 benzo-1,2,5-thiadiazole, benzothiazole, indole, benzotriazole, benzodioxolane, benzodioxane, carbazole, indole and quinazoline.

In the context of aryl and heteroaryl, substituted is  
35 understood as meaning the substitution of the aryl or heteroaryl with R<sup>22</sup>, OR<sup>22</sup>, a halogen, preferably F and/or Cl,

a  $\text{CF}_3$ , a  $\text{CN}$ , an  $\text{NO}_2$ , an  $\text{NR}^{23}\text{R}^{24}$ , a  $\text{C}_{1-6}$ -alkyl (saturated), a  $\text{C}_{1-6}$ -alkoxy, a  $\text{C}_{3-8}$ -cycloalkoxy, a  $\text{C}_{3-8}$ -cycloalkyl or a  $\text{C}_{2-6}$ -alkylene.

- 5 The radical  $\text{R}^{22}$  is H, a  $\text{C}_{1-10}$ -alkyl radical, preferably a  $\text{C}_{1-6}$ -alkyl radical, an aryl or heteroaryl radical or an aryl or heteroaryl radical bonded via a  $\text{C}_{1-3}$ -alkylene group, it being impossible for these aryl and heteroaryl radicals to themselves be substituted by aryl or heteroaryl radicals,
- 10 the radicals  $\text{R}^{23}$  and  $\text{R}^{24}$ , which are identical or different, are H, a  $\text{C}_{1-10}$ -alkyl radical, preferably a  $\text{C}_{1-6}$ -alkyl radical, an aryl or heteroaryl radical or an aryl or heteroaryl radical bonded via a  $\text{C}_{1-3}$ -alkylene group, it being
- 15 impossible for these aryl and heteroaryl radicals to themselves be substituted by aryl or heteroaryl radicals,
- or the radicals  $\text{R}^{23}$  and  $\text{R}^{24}$  together are  $\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2$ ,  $\text{CH}_2\text{CH}_2\text{NR}^{25}\text{CH}_2\text{CH}_2$  or  $(\text{CH}_2)_{3-6}$ , and
- 20 the radical  $\text{R}^{25}$  is H, a  $\text{C}_{1-10}$ -alkyl radical, preferably a  $\text{C}_{1-6}$ -alkyl radical, an aryl or heteroaryl radical or an aryl or heteroaryl radical bonded via a  $\text{C}_{1-3}$ -alkylene group, it being impossible for these aryl and heteroaryl radicals to
- 25 themselves be substituted by aryl or heteroaryl radicals.

'Salt' is understood as meaning any form of the active ingredient according to the invention in which it takes on an ionic (here usually anionic) form, or is charged, and is

30 coupled with a counterion (here usually a cation) or is in solution. This is understood as including complexes of the active ingredient with other molecules and ions, especially complexes that are complexed via ionic interactions.

35 In terms of the present invention, 'physiologically acceptable salts with cations or bases' are understood as

meaning salts of at least one of the compounds according to the invention - usually a (deprotonated) acid - as an anion with at least one, preferably inorganic cation, which are physiologically acceptable, especially when used in humans and/or mammals. Particularly preferred salts are those of alkali metals and alkaline earth metals and also  $\text{NH}_4^+$ , but very particularly preferred salts are monosodium or disodium, monopotassium or dipotassium, magnesium or calcium salts.

In terms of the present invention, 'physiologically acceptable salts with anions or acids' are understood as meaning salts of at least one of the compounds according to the invention - usually protonated, e.g. on the nitrogen - as a cation with at least one anion, which are physiologically acceptable, especially when used in humans and/or mammals. In terms of the present invention, this is understood in particular as meaning the salts formed with physiologically acceptable acids, i.e. salts of the active ingredient in question with inorganic or organic acids, which are physiologically acceptable, especially when used in humans and/or mammals. Examples of physiologically acceptable salts of specific acids are salts of hydrochloric acid, hydrobromic acid, sulfuric acid, methanesulfonic acid, formic acid, acetic acid, oxalic acid, succinic acid, tartaric acid, mandelic acid, fumaric acid, lactic acid, citric acid, glutamic acid, 1,1-dioxo-1,2-dihydro-6-benzo[d]isothiazol-3-one (saccharinic acid), monomethylsebacic acid, 5-oxoproline, hexane-1-sulfonic acid, nicotinic acid, 2-, 3- or 4-aminobenzoic acid, 2,4,6-trimethylbenzoic acid,  $\alpha$ -lipoic acid, acetylglycine, acetylsalicylic acid, hippuric acid and/or aspartic acid. The hydrochloride salt is particularly preferred.

In terms of the present invention, preferred derivatives are the substituted 1,2,3,4-tetrahydroquinoline-2-

carboxylic acid derivatives of formula I in the form of their physiologically acceptable salts with cations or bases. These salts are called 'salts according to the invention' or 'salts according to the invention of substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivatives of formula I' in the description which follows. However, 'salts according to the invention' or 'salts according to the invention of substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivatives of formula I' are not necessarily restricted to physiologically acceptable salts of the substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivatives of formula I with cations or bases, but can optionally also include selected free bases or free acids or physiologically acceptable salts with anions or acids.

One particularly preferred subject of the patent application consists of salts according to the invention of substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivatives of formula I in which  $R^4$  is selected from

H;  $C_1$ - $C_{10}$ -alkyl,  $C_2$ - $C_{10}$ -alkenyl or  $C_2$ - $C_{10}$ -alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted; and  $C_3$ - $C_8$ -cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted; and

$C(O)R^9$ , where  $R^9$  is selected from

H;  $C_1$ - $C_{10}$ -alkyl,  $C_2$ - $C_{10}$ -alkenyl or  $C_2$ - $C_{10}$ -alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted;  $C_3$ - $C_8$ -cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted; and aryl or

heteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted, especially phenethyl, 1-adamantyl, 2-adamantyl, 1-naphthyl or 2-naphthyl, 2-, 3- or 4-pyridyl or thiazolyl.

5

Particularly preferred salts according to the invention of substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivatives of formula I are those in which R<sup>4</sup> is selected from

10

H; C<sub>1</sub>-C<sub>10</sub>-alkyl which is unsubstituted or monosubstituted or polysubstituted; and phenyl which is unsubstituted or monosubstituted or polysubstituted, preferably H, CH<sub>3</sub> or C<sub>2</sub>H<sub>5</sub> and especially H.

15

One preferred subject of the patent application consists of salts according to the invention of substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivatives of formula I in which R<sup>3</sup> is selected from

20

H; C<sub>1</sub>-C<sub>10</sub>-alkyl, C<sub>2</sub>-C<sub>10</sub>-alkenyl or C<sub>2</sub>-C<sub>10</sub>-alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted; C<sub>3</sub>-C<sub>8</sub>-cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted, or a corresponding heterocycle in which at least one ring C atom is replaced by N or O; alkylaryl which is monosubstituted or polysubstituted or unsubstituted; and aryl or heteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted.

25


30

Particularly preferred salts according to the invention of substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivatives of formula I are those in which R<sup>3</sup> is selected from

35

H; C<sub>1</sub>-C<sub>4</sub>-alkyl which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted; and phenyl, benzyl or phenethyl which is  
 5 monosubstituted or polysubstituted or unsubstituted, preferably H, CH<sub>3</sub> or C<sub>2</sub>H<sub>5</sub> and especially H.

Particularly preferred salts according to the invention of substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid  
 10 derivatives of formula I are those in which R<sup>1</sup> and R<sup>2</sup> together form -O-CH<sub>2</sub>-CH<sub>2</sub>-, (-CH<sub>2</sub>-)<sub>n</sub> where n = 3-6, preferably 3 or 6, -CH=CH-CH<sub>2</sub>-, -CH=CH-CH<sub>2</sub>-CH<sub>2</sub>-,



15 preferably -CH=CH-CH<sub>2</sub>- or -CH=CH-CH<sub>2</sub>-CH<sub>2</sub>- and especially -CH=CH-CH<sub>2</sub>-.

Another preferred subject consists of salts according to  
 20 the invention of substituted 1,2,3,4-tetraquinoline-2-carboxylic acid derivatives of formula I in which R<sup>1</sup> is selected from

phenyl, naphthyl or anthracenyl which is unsubstituted  
 25 or monosubstituted or polysubstituted; and OR<sup>19</sup> or SR<sup>19</sup>, where R<sup>19</sup> is selected from

C<sub>1</sub>-C<sub>6</sub>-alkyl which is branched or unbranched and monosubstituted or polysubstituted or

unsubstituted; C<sub>3</sub>-C<sub>8</sub>-cycloalkyl which is saturated  
or unsaturated and monosubstituted or  
polysubstituted or unsubstituted; and aryl which  
is monosubstituted or polysubstituted or  
5 unsubstituted;

preferably anthracenyl, naphthyl or, in particular,  
phenyl which is unsubstituted or monosubstituted or  
polysubstituted by a substituent selected from

10 F; Cl; Br; I; methoxy; ethoxy; propoxy; methyl;  
ethyl; propyl (n-propyl, i-propyl); butyl  
(n-butyl, i-butyl, t-butyl); carboxyl; nitro;  
benzyloxy; phenyl; hydroxyl; phenoxy;  
15 trifluoromethyl; dioxolyl and SCH<sub>3</sub>;

or OR<sup>19</sup> or SR<sup>19</sup>, where R<sup>19</sup> is selected from

C<sub>1</sub>-C<sub>4</sub>-alkyl which is branched or unbranched and  
20 monosubstituted or polysubstituted or  
unsubstituted; C<sub>3</sub>-C<sub>8</sub>-cycloalkyl which is saturated  
or unsaturated and monosubstituted or  
polysubstituted or unsubstituted; and aryl which  
is monosubstituted or polysubstituted or  
25 unsubstituted;

especially unsubstituted phenyl, naphthyl and  
anthracenyl, O-hydroxyethyl, ethoxynaphthyl, 4-  
hydroxy-3-methoxyphenyl, 4-propoxyphenyl, 2,3,4-  
30 trimethylphenyl, 2,4,5-trimethoxyphenyl, SCH<sub>3</sub>, 2-  
chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 2-  
fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 2-  
bromophenyl, 3-bromophenyl, 4-bromophenyl, 2,6-  
dichlorophenyl, 4-carboxyphenyl, 3-nitrophenyl, 2,4,6-  
35 trimethylphenyl, 2,5-dimethylphenyl, 3,4-  
dimethoxyphenyl, 4-benzyloxy-3-methoxyphenyl, 3-



5 methylphenyl, 4-methoxyphenyl, 4-biphenyl, 4-methylphenyl, 4-ethoxyphenyl, 2-methylphenyl, 2,4-dimethylphenyl, 2,6-dimethylphenyl, 4-hydroxy-3-methoxyphenyl, 4-methylhydroxyphenyl, 4-hydroxyphenyl, 4-phenoxyphenyl, 4-nitrophenyl, 4-chloromethylphenyl, 4-tert-butylphenyl, 3,5-bis(trifluoromethyl)phenyl, 4-acetoxyphenyl, 4-cyanophenyl, 2-methoxyphenyl, 2,6-difluorophenyl, 2-trifluoromethylphenyl, 3-trifluoromethylphenyl, 4-trifluoromethylphenyl, 3-methoxyphenyl, 2-, 3- or 4-benzyloxyphenyl, S-phenyl or 6-chlorobenzo[1,3]dioxol-5-yl.

Another preferred subject of the patent application consists of salts according to the invention of substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivatives of formula I in which R<sup>2</sup> is selected from

20 H; C<sub>1</sub>-C<sub>4</sub>-alkyl which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted; and phenyl which is monosubstituted or polysubstituted or unsubstituted, preferably H, unsubstituted phenyl, 4-methoxyphenyl or CH<sub>3</sub> and especially H.

One preferred subject of the patent application consists of salts according to the invention of substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivatives of formula I in which R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> independently of one another are selected from

30 H; F; Cl; Br; I; CN; NO<sub>2</sub>; and C<sub>1</sub>-C<sub>10</sub>-alkyl, C<sub>2</sub>-C<sub>10</sub>-alkenyl or C<sub>2</sub>-C<sub>10</sub>-alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted;

35 OR<sup>14</sup>, C(O)R<sup>14</sup>, C(O)OR<sup>14</sup> or SR<sup>14</sup>, R<sup>14</sup> being selected from

H; C<sub>1</sub>-C<sub>10</sub>-alkyl, C<sub>2</sub>-C<sub>10</sub>-alkenyl or C<sub>2</sub>-C<sub>10</sub>-alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted; C<sub>3</sub>-C<sub>8</sub>-cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted, or a corresponding heterocycle in which at least one ring C atom is replaced by S, O or N; alkylaryl or alkylheteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted; and aryl or heteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted; and

NR<sup>15</sup>R<sup>16</sup> or NR<sup>15</sup>C(O)R<sup>16</sup>, R<sup>15</sup> and R<sup>16</sup> independently of one another being selected from

H; O; C<sub>1</sub>-C<sub>10</sub>-alkyl, C<sub>2</sub>-C<sub>10</sub>-alkenyl or C<sub>2</sub>-C<sub>10</sub>-alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted.

Particularly preferred salts according to the invention of substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivatives of formula I are those in which R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> independently of one another are selected from

H; F; Cl; Br; I; CN; NO<sub>2</sub>; and C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl or C<sub>2</sub>-C<sub>6</sub>-alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted;

OR<sup>14</sup>, C(O)R<sup>14</sup>, C(O)OR<sup>14</sup> or SR<sup>14</sup>, R<sup>14</sup> being selected from

H; C<sub>1</sub>-C<sub>4</sub>-alkyl which is branched or unbranched and monosubstituted or polysubstituted or

unsubstituted; and aryl which is monosubstituted or polysubstituted or unsubstituted; preferably, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> independently of one another are selected from

5

H; F; Cl; Br; I; CN; and C<sub>1</sub>-C<sub>4</sub>-alkyl which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted;

10

OR<sup>14</sup> or SR<sup>14</sup>, R<sup>14</sup> being selected from

15

C<sub>1</sub>-C<sub>4</sub>-alkyl which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted; and aryl which is monosubstituted or polysubstituted or unsubstituted;

in particular, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> independently of one another are selected from

20

H; F; Cl; Br; I; CN; CH<sub>3</sub>; CF<sub>3</sub>; t-butyl; i-butyl; -OCH<sub>3</sub>; -OCF<sub>3</sub>; -SCH<sub>3</sub> and -O-phenyl.

Very particularly preferred salts according to the invention of substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivatives of formula I are those in which

25

R<sup>5</sup>, R<sup>6</sup> and R<sup>8</sup> are H and R<sup>7</sup> is Cl, or  
R<sup>5</sup> and R<sup>7</sup> are H and R<sup>6</sup> and R<sup>8</sup> are Cl.

Particularly preferred subjects consist of the salts according to the invention of the following substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivatives:

30

7,9-dichloro-3a,4,5,9b-tetrahydro-3H-cyclopenta[c]-quinoline-4-carboxylic acid,

35

8-chloro-3a,4,5,9b-tetrahydro-3H-cyclopenta[c]quinoline-4-carboxylic acid,

6-chloro-7-trifluoromethyl-4-(2,4,6-trimethylphenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

5

4-(2-hydroxyethoxy)-6-trifluoromethoxy-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

6-iodo-4-(4-methoxyphenyl)-3-methyl-1,2,3,4-

10 

tetrahydroquinoline-2-carboxylic acid,

5,7-dichloro-4-phenyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

15 

5,7-dichloro-4-m-tolyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

5,7-dichloro-4-p-tolyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

20

5,7-dichloro-4-(2,4-dimethylphenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

5,7-dichloro-4-(2-fluorophenyl)-1,2,3,4-

25 

tetrahydroquinoline-2-carboxylic acid,

5,7-dichloro-4-(3-fluorophenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

30 

5,7-dichloro-4-(4-fluorophenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

5,7-dichloro-4-(2-chlorophenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

35

5,7-dichloro-4-(4-chlorophenyl)-1,2,3,4-

tetrahydroquinoline-2-carboxylic acid,

5,7-dichloro-4-(3-bromophenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

5

5,7-dichloro-4-(4-bromophenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

7,8-dichloro-4-(2-chlorophenyl)-1,2,3,4-

10 tetrahydroquinoline-2-carboxylic acid,

6-cyano-4-(2,3,4-trimethoxyphenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

15 6,8,9-trichloro-2,3,3a,4,5,9b-hexahydrofuro[3,2-c]-quinoline-4-carboxylic acid,

8-methoxy-4-(4-methoxyphenyl)-3-methyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

20

5,6,8-trichloro-4-(4-hydroxyphenyl)-3-methyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

4-(3,4-dimethoxyphenyl)-8-iodo-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

25

6-iodo-4-(4-methylsulfanylphenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

4-(4-ethoxy-3-methoxyphenyl)-6-phenoxy-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

30

4-(2-ethoxynaphthalen-1-yl)-6-iodo-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

35

8-chloro-4-(4-propoxyphenyl)-1,2,3,4-tetrahydroquinoline-2-

carboxylic acid,

4-(2,4-dimethoxy-3-methylphenyl)-6-phenoxy-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

5

4-anthracen-9-yl-6-chloro-8-methyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

6-sec-butyl-4-naphthalen-1-yl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

10

4-(4-hydroxyphenyl)-3-methyl-8-phenoxy-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

8-chloro-6-fluoro-4-naphthalen-2-yl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

15

4-(4-methoxyphenyl)-3-methyl-6-phenoxy-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

20

6-chloro-8-fluoro-4-m-tolyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

8-chloro-6-fluoro-4-m-tolyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

25

4-(4-bromophenyl)-6-chloro-8-fluoro-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

7,8-dichloro-4-(2,4-dimethylphenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

30

6-chloro-4-(4-chlorophenyl)-7-trifluoromethyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

35

4-(2-chlorophenyl)-6-cyano-1,2,3,4-tetrahydroquinoline-2-

carboxylic acid,

6-bromo-8-chloro-4-(2,4-dimethylphenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

5

6-bromo-4-(2-bromophenyl)-8-chloro-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

10 4-(4-hydroxy-3-methoxyphenyl)-3-methyl-6-methylsulfanyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

6-cyano-3,4-bis(4-methoxyphenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

15 5,7-dichloro-4-(4-chlorophenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

5,7-dichloro-4-(3-chlorophenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

20

5,7-dichloro-4-(4-chlorophenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

25 1,3-dichloro-5,6,6a,7,8,12b-hexahydrobenzo[k]-phenanthridine-6-carboxylic acid,

1,3-dichloro-5,6a,7,11b-tetrahydro-6H-indeno[2,1-c]-quinoline-6-carboxylic acid,

30 5,7-dichloro-4-(3,5-dimethylphenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid and

7,9-dichloro-2,3,3a,4,5,9b-hexahydro-1H-cyclopenta[c]-quinoline-4-carboxylic acid,

35

particular preference being given to sodium 7,9-

dichloro-3a,4,5,9b-tetrahydro-3H-cyclopenta[c]-  
quinoline-4-carboxylate or sodium 7,9-dichloro-  
2,3,3a,4,5,9b-hexahydro-1H-cyclopenta[c]quinoline-4-  
carboxylate, especially sodium 7,9-dichloro-3a,4,5,9b-  
5 tetrahydro-3H-cyclopenta[c]quinoline-4-carboxylate.

Particularly preferred salts according to the invention of  
substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid  
derivatives of formula I are the alkali metal salts,  
10 preferably the sodium or potassium salts and especially the  
sodium salts.

The invention also provides processes for the preparation  
of salts according to the invention of a substituted  
15 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivative.

Various processes for the preparation of  
tetrahydroquinolines are described in the literature:

- a solid phase preparation (WO 98/34111),
- 20 • multistage procedures (WO 98/42673; Bioorganic and  
Medicinal Chemistry Letters, vol. 2, p. 371, 1992;  
Journal of Heterocyclic Chemistry, vol. 25, p. 1831,  
1988; Journal of the Chemical Society, Perkin  
Transactions I (1989), page 2245) or
- 25 • a Lewis acid-catalyzed "one-pot" process (Journal of the  
Chemical Society, Chemical Communications, 1999, p. 651;  
Journal of the American Chemical Society, vol. 118, p.  
8977, 1996).


30 However, all these processes clearly have some  
disadvantages.

In contrast to these processes, the so-called basic process  
described here is a process mediated by trifluoroacetic  
35 acid - preferably a "one-pot" process - in which one  
aromatic amine component, one aldehyde component and one



electron-rich olefin component are reacted with one another.

In the basic process, substituted 1,2,3,4-  
 5 tetrahydroquinoline-2-carboxylic acid derivatives of  
 formula I in which  $R^4 = H$ , the other radicals having one of  
 the meanings already mentioned, are prepared first.  
 Anilines of formula II, in which  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  each  
 10 independently of one another have one of the meanings  
 already indicated or are provided with a protecting group,



glyoxalic acid esters or optionally glyoxalic acid of  
 15 formula III and olefins of formula IV, in which  $R^1$ ,  $R^2$  and  
 $R^3$  each independently of one another have one of the  
 meanings already indicated or are provided with a  
 protecting group, are reacted with trifluoroacetic acid at  
 between 0°C and 100°C. The reaction time is preferably  
 20 0.25 - 12 h and particularly preferably at most 2 h, the  
 reaction is carried out preferably at a temperature of  
 between 20 and 40°C and particularly preferably at room  
 temperature, and/or the reaction is a one-pot reaction.  
 When this basic process has ended, any existing ester  
 25 groups can be saponified and/or the product formed in the  
 basic process can optionally be brought into contact with a  
 strong base, which may already contain the desired cation,  
 in order to form a salt.

One decisive advantage of the process according to the invention is that it produces the desired systems according to a domino reaction (imine formation and subsequent aza Diels-Alder reaction) with very high selectivities coupled  
5 with good yields.

Without having to carry out a linking or cleaving step as in the case of the solid phase preparation, and also without purification of the intermediates as in the case of  
10 the solution chemistry described, the process according to the invention is distinguished not only by its ease of implementation but also by its purification method. Washing several times with non-polar solvents, for example n-hexane, makes it possible for the most part to obtain the  
15 products in high purity. In other cases, they can be purified by means of column chromatography. In particular, the compounds of formula I can be obtained as the pure diastereoisomers by the washing processes with non-polar solvents, for example n-hexane, or by crystallization of  
20 their salts.

In general, in one advantageous embodiment of the preparative process, when the formation of a compound of formula I has ended, the compound is brought into contact  
25 with a strong base, which may already contain the desired cation, and the resulting salt according to the invention is then purified.

The majority of the reactants used here, especially those  
30 of formulae II, III and IV, are commercially available or can be prepared by simple synthesis steps known to those skilled in the art.

In consecutive reactions following the basic process, the  
35 products formed in the basic process can be converted to consecutive products of formula I according to the

invention by a procedure known to those skilled in the art, whereby initially the hydrogen on  $R^4$  is substituted.

Thus, when the basic reaction has ended, if the product is  
5 to be a substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivative of formula I where  $R^4$  = alkylformyl, acyl, sulfenyl or sulfonyl, the reaction product can be reacted with the appropriate chloroformate or fluoroformate, acid chloride, sulfenyl chloride or  
10 sulfonyl chloride in the presence of a base, preferably triethylamine, pyridine or NaOH, in water, a dioxane/water mixture or a THF/water mixture, at a temperature of between 0 and 20°C (J. Org. Chem., 1989, 54, 5574-5580).

15 Likewise, when the basic reaction has ended, if the product is to be a substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivative of formula I where  $R^4$  =  $C(S)NR^{11}R^{12}$ , the reaction product can be reacted with a thionating reagent, preferably Lawesson's reagent (2,4-bis(4-methoxyphenyl)-2,4-dithioxo-1,3,2,4-dithiaphosphetane), in an organic solvent, preferably THF  
20 or toluene, at a temperature of 30-50°C.

Alternatively, when the basic reaction has ended, if the  
25 product is to be a substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivative of formula I where  $R^4$  =  $C(O)NR^{11}R^{12}$  or  $C(S)NR^{11}R^{12}$ , the reaction product can be reacted with potassium cyanate or potassium isothiocyanate in water at temperatures of up to 100°C, or with an organic  
30 isocyanate or isothiocyanate in an alcohol, preferably methanol, ethanol or isopropanol, at temperatures up to the boiling point.

Again, when the basic reaction has ended, if the product is  
35 to be a substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivative of formula I where  $R^4$  =

$C(NR^{11})NR^{12}R^{13}$ , the reaction product can be reacted under alkaline conditions with an O-methylisourea or S-methylisothiourea at temperatures of 20-50°C, preferably ethanolic or methanolic NaOH or KOH.

5

Yet again, when the basic reaction has ended, if the product is to be a substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivative of formula I where  $R^4 = C(O)NR^{11}R^{12}$ , the reaction product can be reacted with propanone-2-semicarbazone in water/glacial acetic acid at 30-60°C.

10

Likewise, when the basic reaction has ended, if the product is to be a substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivative of formula I where  $R^4 = C(S)NR^{11}R^{12}$ , the reaction product can be reacted with  $CS_2$  and a hydrazine in water/NaOH at 30-60°C.

15

As the final possibility to be mentioned here, when the basic reaction has ended, if the product is to be a substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivative of formula I where  $R^4 =$  alkyl, benzyl or phenethyl, the reaction product can be reacted with an appropriate alkylating halide, benzyl halide or phenethyl halide and a suitable base, preferably sodium hydride or potassium tert-butyrate, in a solvent, for example ethanol, at between 0 and 100°C (J. Org. Chem., 1947, 12, 760; Zh. Obshch. Khim., 1942, 12, 418).

20

25

30

35

Under many of said reaction conditions, OH, SH and  $NH_2$  groups may possibly enter into unwanted secondary reactions. It is therefore preferable to provide them with protecting groups or, in the case of  $NH_2$ , to replace it with  $NO_2$ , and to cleave the protecting group, or reduce the  $NO_2$  group, before purifying the end product. The patent application therefore also provides a modification of the

process described above wherein, in the starting compounds, at least one OH group has been replaced by an  $\text{OSi(Ph)}_2\text{tert-butyl}$  group, at least one SH group has been replaced by an S-p-methoxybenzyl group and/or at least one  $\text{NH}_2$  group has  
5 been replaced by an  $\text{NO}_2$  group, and at least one and preferably all of the  $\text{OSi(Ph)}_2\text{tert-butyl}$  groups are cleaved with tetrabutylammonium fluoride in tetrahydrofuran, and/or at least one and preferably all of the p-methoxybenzyl groups are cleaved with a metal amine, preferably sodamine,  
10 and/or at least one and preferably all of the  $\text{NO}_2$  groups are reduced to  $\text{NH}_2$  before the end product is purified.

Furthermore, carboxylic acid or thiocarboxylic acid groups are sometimes unstable under said reaction conditions, so  
15 it is preferable to use their methyl esters in the reactions and then to saponify the product of the process with KOH solution or NaOH solution in methanol at  $40^\circ\text{C}$  -  $60^\circ\text{C}$ . The invention therefore also provides a modification of the processes described above wherein, before the end  
20 product is purified, a product of the process having at least one  $\text{C(O)OCH}_3$ ,  $\text{OC(O)OCH}_3$  and/or  $\text{C(S)OCH}_3$  group is saponified with KOH solution or NaOH solution in methanol or ethanol at  $0^\circ\text{C}$  -  $100^\circ\text{C}$ , preferably at  $40^\circ\text{C}$  -  $60^\circ\text{C}$ .

25 Therefore, it can also be favourable, for the preparation of (salts according to the invention of) substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivatives of formula I in which  $\text{R}^3 = \text{H}$ , to use for the basic process starting materials of formula III in which  $\text{R}^3 \neq \text{H}$  and is  
30 preferably alkyl, especially  $\text{CH}_3$  and  $\text{C}_2\text{H}_5$ . After the basic process and also the consecutive reactions that may follow it, the reaction product is saponified with an appropriate base, preferably NaOH (for example 6 N) or KOH, in ethanol or methanol at temperatures of between 0 and  $100^\circ\text{C}$ ,  
35 preferably of  $40^\circ\text{C}$  -  $60^\circ\text{C}$  (Organikum, 1990, p. 418).

The following procedure is used to prepare the salts, especially the physiologically acceptable salts with cations or bases:

- 5 One equivalent of a compound of formula I, preferably an imino acid or a carboxylic acid, particularly where  $R^3 = H$ , is suspended in a small volume of water and one equivalent of 1 N aqueous alkaline solution, for example NaOH or KOH, is added. If the solubility is poor, methanol is added  
10 dropwise until the compound has completely dissolved. After stirring at room temperature, the solution is concentrated on a rotary evaporator and the residual solution is frozen at low temperatures in an isopropanol/dry ice mixture and freeze-dried. The salts, especially of  
15 imino acids or carboxylic acids and preferably the sodium or potassium salts, are usually obtained as colourless solids.

- An alternative possibility is to prepare the potassium or  
20 sodium salts with potassium or sodium trimethylsilanolate (E.D. Laganis, B.L. Chenard, Tetrahedron Letters, 25, 5831-5834 (1984)). Potassium or sodium trimethylsilanolate is dissolved under nitrogen in an organic solvent (e.g. dichloromethane, toluene, THF) and the ester or acid is  
25 added all at once. The reaction mixture is stirred for several hours at room temperature and filtered. The usually colourless solid is washed and dried under vacuum. The potassium or sodium salts are obtained as solids.

- 30 The salts according to the invention of substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivatives are toxicologically safe, so they are suitable as pharmaceutical active ingredients in drugs.

- 35 The invention therefore also provides a drug containing, as the active ingredient, at least one salt according to the invention of a substituted 1,2,3,4-tetrahydroquinoline-2-

carboxylic acid derivative of formula I in the form shown, or in the form of the acid or base, or in the form of its salts, especially the physiologically acceptable salts, or in the form of its solvates, especially the hydrates, particularly in the form of its physiologically acceptable salts with cations or bases or with anions or acids, optionally in the form of its racemates or its pure stereoisomers, especially enantiomers or diastereoisomers, or in the form of mixtures of the stereoisomers, especially the enantiomers or diastereoisomers, in any desired mixing ratio, and optionally containing suitable additives and/or auxiliary substances and/or optionally other active ingredients.

The drugs according to the invention can be administered as liquid dosage forms, i.e. injection solutions, drops or juices, or as semisolid dosage forms, i.e. granules, tablets, pellets, patches, capsules, plasters or aerosols, and optionally contain, apart from at least one salt according to the invention of a substituted tetrahydroquinoline derivative, excipients, fillers, solvents, diluents, colorants and/or binders, depending on the galenical form. The choice of auxiliary substances and the amounts thereof to be used depend on whether the drug is to be administered orally, perorally, parenterally, intravenously, intraperitoneally, intradermally, intramuscularly, intranasally, buccally, rectally or locally, for example to infections on the skin, mucous membranes or eyes. Suitable preparations for oral administration are those in the form of tablets, dragees, capsules, granules, drops, juices and syrups, and suitable preparations for parenteral, topical and inhalational administration are solutions, suspensions, readily reconstitutable dry preparations and sprays. Salts according to the invention of substituted tetrahydroquinoline derivatives in a depot in dissolved form or in a plaster, optionally with the addition of skin

penetration promoters, are suitable preparations for transdermal application. Preparative forms for oral or transdermal administration can release the salts according to the invention of substituted tetrahydroquinoline derivatives in a delayed manner. The amount of active ingredient to be administered to the patient varies according to the patient's weight, the mode of administration, the indication and the severity of the disease. Conventionally, 2 to 500 mg/kg of at least one salt according to the invention of a substituted tetrahydroquinoline derivative of formula I are administered.

Preferably, the salts according to the invention of substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivatives are used for the treatment of pain, especially chronic and neuropathic pain, as well as migraine, so the invention also provides the use of at least one salt according to the invention of a substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivative of formula I, optionally in the form of its racemates or its pure stereoisomers, especially enantiomers or diastereoisomers, or in the form of mixtures of the stereoisomers, especially the enantiomers or diastereoisomers, in any desired mixing ratio, for the preparation of a drug for the treatment of pain, especially neuropathic and/or chronic pain, and/or for the treatment of migraine.

The affinity for the NMDA receptor gives rise to other areas of application since it is known that NMDA antagonists have *inter alia* a neuroprotective action and can therefore also be used satisfactorily for syndromes associated with neurodegeneration and neural damage, such as Parkinson's disease and Huntington's chorea, etc. Other indications of the NMDA antagonists according to the invention are epilepsy, glaucoma, osteoporosis,



ototoxicity, withdrawal symptoms associated with alcohol and/or drug abuse, stroke, including related cerebral ischaemia, cerebral infarction, cerebral oedema, hypoxia and anoxia, and also use for anxiolysis and in anaesthesia.

5 The invention therefore also provides the use of at least one salt according to the invention of a substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivative of formula I, optionally in the form of its racemates or its pure stereoisomers, especially enantiomers or  
10 diastereoisomers, or in the form of mixtures of the stereoisomers, especially the enantiomers or diastereoisomers, in any desired mixing ratio, for the preparation of a drug for the treatment/prophylaxis of epilepsy, Parkinson's disease, Huntington's chorea,  
15 glaucoma, ototoxicity, withdrawal symptoms associated with alcohol and/or drug abuse, stroke, cerebral ischaemia, cerebral infarction, cerebral oedema, hypoxia and anoxia, and/or for anxiolysis and/or anaesthesia.

20 Surprisingly, it has been found that the substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivatives according to the invention are also very suitable for other indications and especially for the treatment of urinary incontinence, pruritus, tinnitus aurium and/or diarrhoea.

25 The patent application therefore also provides the use of at least one salt according to the invention of a substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivative of formula I, optionally in the form of its racemates or its pure stereoisomers, especially enantiomers  
30 or diastereoisomers, or in the form of mixtures of the stereoisomers, especially the enantiomers or diastereoisomers, in any desired mixing ratio, for the preparation of a drug for the treatment of urinary incontinence, pruritus, tinnitus aurium and/or diarrhoea.

35 However, the compounds according to the invention are also

effective in other indications. The patent application therefore also provides the use of at least one salt according to the invention of a substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivative of formula I, optionally in the form of its racemates or its pure stereoisomers, especially enantiomers or diastereoisomers, or in the form of mixtures of the stereoisomers, especially the enantiomers or diastereoisomers, in any desired mixing ratio, for the treatment/prophylaxis of schizophrenia, Alzheimer's disease, psychosis due to a raised amino acid level, AIDS-related dementia, encephalomyelitis, Tourette syndrome, perinatal asphyxia, inflammatory and allergic reactions, depression, drug and/or alcohol abuse, gastritis, diabetes, cardiovascular disease, respiratory tract disease, cough and/or mental disease.

The invention also provides a method of treating a non-human mammal or a human in need of a treatment for medically relevant symptoms, by the administration of a therapeutically effective dose of a salt according to the invention of a substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivative of formula I, optionally in the form of its racemates or its pure stereoisomers, especially enantiomers or diastereoisomers, or in the form of mixtures of the stereoisomers, especially the enantiomers or diastereoisomers, in any desired mixing ratio, or a drug according to the invention. The invention relates especially to appropriate methods for the treatment of pain, especially neuropathic and/or chronic pain, and/or for the treatment of migraine, for the treatment of urinary incontinence, pruritus, tinnitus aurium and/or diarrhoea, for the treatment/prophylaxis of epilepsy, Parkinson's disease, Huntington's chorea, glaucoma, osteoporosis, ototoxicity, withdrawal symptoms associated with alcohol and/or drug abuse, stroke, cerebral ischaemia, cerebral infarction, cerebral oedema, hypoxia and anoxia, and/or for

anxiolysis and/or anaesthesia, or for the treatment/  
prophylaxis of schizophrenia, Alzheimer's disease,  
psychosis due to a raised amino acid level, AIDS-related  
dementia, encephalomyelitis, Tourette syndrome, perinatal  
5 asphyxia, inflammatory and allergic reactions, depression,  
drug and/or alcohol abuse, gastritis, diabetes,  
cardiovascular disease, respiratory tract disease, cough  
and/or mental disease.

10 The invention is further illustrated below by means of  
Examples, which do not imply a limitation.

### **Examples**

15 The Examples which follow show compounds according to the  
invention, their preparation and efficacy tests performed  
therewith.

The following information is generally applicable:

20

The chemicals and solvents used were acquired commercially  
from the conventional suppliers (Acros, Avocado, Aldrich,  
Fluka, Lancaster, Maybridge, Merck, Sigma, TCI, etc.) or  
synthesized.

25

In particular, some of the compounds used are synthesized  
as synthetic moieties by known procedures before the basic  
synthesis described below.

30 The thin layer chromatography tests were carried out with  
HPTLC precoated plates, silica gel 60 F 254, from E. Merck,  
Darmstadt.

The yields of the compounds prepared are not optimized.

35

The analysis was performed by ESI mass spectroscopy.

The compounds are numbered, the data in brackets corresponding in principle to the number of the assigned compound.

5

**Example 0****Basic process for the preparation of the basic compounds of formula I**

10

a) One equivalent of aniline derivative and one equivalent of trifluoroacetic acid are dissolved in 6 ml/mmol of acetonitrile at room temperature, with stirring, and 1.1 equivalents of ethyl glyoxalate (50% in toluene) or 1.1 equivalents of glyoxalic acid monohydrate are then added. After ten minutes, 3 equivalents of the olefin component are added and the course of the reaction is followed by thin layer chromatography (solvent system: diethyl ether/hexane, 1:1). The reaction has ended after 2 hours (TLC check). An excess of saturated aqueous sodium hydrogencarbonate solution is added to the reaction mixture and the organic phase is extracted three times with diethyl ether. The organic phase is washed with water until the washings are neutral and dried over magnesium sulfate, the magnesium sulfate is filtered off and washed with diethyl ether, the product phase is concentrated and the product is isolated by recrystallization or silica gel chromatography. The 1,2,3,4-tetrahydroquinoline-2-carboxylic acid ester is characterized by ESI mass spectrometry.

30

b) Optional subsequent preparation of the free 1,2,3,4-tetrahydroquinoline-2-carboxylic acids

The above-described 1,2,3,4-tetrahydroquinoline-2-carboxylic acid ester (1 equivalent) is dissolved in 4 ml/mmol of ethanol, and 1.2 equivalents of 6 N aqueous

35

sodium hydroxide solution are added at room temperature, with stirring. The course of the ester saponification is followed by thin layer chromatography (solvent system: diethyl ether/hexane, 1:1) and has ended after 30 minutes (TLC check). The reaction mixture is concentrated on a rotary evaporator, taken up in approx. 10 ml of water and adjusted to pH 1 with 32% HCl. The aqueous solution is extracted five times with diethyl ether, dried over magnesium sulfate and concentrated.

#### Automated process

A round-bottom screw-threaded glass tube (diameter 16 mm, length 125 mm) was provided with a stirrer and sealed with a screw cap comprising a septum. The tube was placed in a stirring block thermostatted at 20°C. The following reactants were then pipetted successively into the tube:

1 ml of a solution of 0.1 M trifluoroacetic acid and 0.1 M aniline component in acetonitrile;

1 ml of a 0.11 M solution of the aldehyde in acetonitrile;

1 ml of a 0.3 M solution of the olefin in acetonitrile.

The reaction mixture was stirred for 10 h at 20°C in one of the stirring blocks. The reaction solution was then filtered, the tube being rinsed with twice 1.5 ml of 7.5% NaHCO<sub>3</sub> solution.

2 ml of ethyl acetate were added to the reaction mixture on a vortex shaker and the mixture was shaken. It was centrifuged briefly to form the phase boundary. This was detected optically and the organic phase was pipetted off. In the next step a further 2 ml of ethyl acetate were added to the aqueous phase, the mixture was shaken and centrifuged and the organic phase was pipetted off. The combined organic phases were dried over 2.4 g of MgSO<sub>4</sub> (granulated). The solvent was stripped off in a vacuum

centrifuge.

The free 1,2,3,4-tetrahydroquinoline-2-carboxylic acid or the ester is characterized by ESI mass spectrometry.

5

In principle, in the case of compounds where  $R^3 \neq H$ , both the automated process and the normal basic process can be followed by a saponification using methods known to those skilled in the art, for example with KOH solution or NaOH solution in methanol or ethanol at 0°C - 100°C, preferably at 40°C - 60°C.


Examples 1 to ... which now follow show the preparation of the basic compounds of formula I by one of the processes of Example 0, salts according to the invention of said compounds being prepared after the particular preparation described.

15

#### Example 1

20

7,9-Dichloro-3a,4,5,9b-tetrahydro-3H-cyclopenta[c]-quinoline-4-carboxylic acid (**1**)



25

Compound **1** was saponified with 1.0 ml of sodium hydroxide solution (6 N, water) in 20.0 ml of ethanol. The ethanolic solution was concentrated on a rotary evaporator, the residue was taken up in water, 6 N HCl was added and the

aqueous solution was extracted three times with ether. The organic phase was washed with water until the washings were neutral, dried over magnesium sulfate and concentrated on a rotary evaporator.


5 An ESI-MS was run to characterize the product:

MS (EI) m/z: 284 (M\*)

### Example 2

10

8-Chloro-3a,4,5,9b-tetrahydro-3H-cyclopenta[c]quinoline-4-carboxylic acid (**2**)



15


Compound **2** was saponified with 1.0 ml of sodium hydroxide solution (6 N, water) in 20.0 ml of ethanol. The ethanolic solution was concentrated on a rotary evaporator, the residue was taken up in water, 6 N HCl was added and the aqueous solution was extracted three times with ether. The organic phase was washed with water until the washings were neutral, dried over magnesium sulfate and concentrated on a rotary evaporator.

20

An ESI-MS was run to characterize the product:

25

MS (EI) m/z: 250 (M\*)

**Example 3**


- 5 6-Chloro-7-trifluoromethyl-4-(2,4,6-trimethylphenyl)-  
1,2,3,4-tetrahydroquinoline-2-carboxylic acid (**3**)

Compound **3** was prepared by the basic process from 5.0 mmol  
of 4-chloro-4-trifluoromethylaniline, 5.5 mmol of glyoxalic  
10 acid monohydrate and 15.0 mmol of 2,4,6-trimethylstyrene in  
30 ml of acetonitrile.

An ESI-MS was run to characterize the product:


15 MS (EI) m/z: 398.1 (M\*)



**Example 4**


- 5 4-(2-Hydroxyethoxy)-6-(trifluoromethoxy)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid **(4)**

Compound **4** was prepared by the automated process from 4-(trifluoromethoxy)aniline, glyoxylic acid and ethylene  
10 glycol monovinyl ether.

**Example 5**

- 15 6-Iodo-4-(4-methoxyphenyl)-3-methyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid **(5)**

Compound **5** was prepared by the automated process from 4-iodoaniline, glyoxylic acid and trans-anethole.  
20


**Example 6**

- 5 5,7-Dichloro-4-phenyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid (**6**)

Compound **6** was prepared by the basic process from 5.0 mmol of 3,5-dichloroaniline, 5.5 mmol of ethyl glyoxalate solution (50%, toluene), 15.0 mmol of styrene and 5.0 mmol of trifluoroacetic acid in 30.0 ml of acetonitrile. The subsequent saponification was carried out with 1.0 ml of sodium hydroxide solution (6 N, water) in 20.0 ml of ethanol.

- 15 An ESI-MS was run to characterize the product:

MS (EI) m/z: (M\*) 315

**Example 7**

20

5,7-Dichloro-4-*m*-tolyl-1,2,3,4-tetrahydroquinoline-2-

carboxylic acid (7)


Compound **7** was prepared by the basic process from 5.0 mmol of 3,5-dichloroaniline, 5.5 mmol of ethyl glyoxalate solution (50%, toluene), 15.0 mmol of 3-methylstyrene and 5.0 mmol of trifluoroacetic acid in 30.0 ml of acetonitrile. The subsequent saponification was carried out with 1.0 ml of sodium hydroxide solution (6 N, water) in 20.0 ml of ethanol.

10 An ESI-MS was run to characterize the product:

MS (EI) m/z: (M<sup>+</sup>) 335

### Example 8

15



5,7-Dichloro-4-*p*-tolyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid (**8**)

20


Compound **8** was prepared by the basic process from 5.0 mmol of 3,5-dichloroaniline, 5.5 mmol of ethyl glyoxalate solution (50%, toluene), 15.0 mmol of 4-methylstyrene and 5.0 mmol of trifluoroacetic acid in 30.0 ml of acetonitrile. The subsequent saponification was carried out with 1.0 ml of sodium hydroxide solution (6 N, water) in 20.0 ml of ethanol.

An ESI-MS was run to characterize the product:

MS (EI) m/z: (M\*) 335

**Example 9**

5



5,7-Dichloro-4-(2,4-dimethylphenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid (**9**)

10


Compound **9** was prepared by the basic process from 5.0 mmol of 3,5-dichloroaniline, 5.5 mmol of ethyl glyoxalate solution (50%, toluene), 15.0 mmol of 2,4-dimethylstyrene and 5.0 mmol of trifluoroacetic acid in 30.0 ml of acetonitrile. The subsequent saponification was carried out with 1.0 ml of sodium hydroxide solution (6 N, water) in 20.0 ml of ethanol.

15

An ESI-MS was run to characterize the product:

20

MS (EI) m/z: (M\*) 349

**Example 10**

- 5 5,7-Dichloro-4-(2-fluorophenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid (**10**)


Compound **10** was prepared by the basic process from 5.0 mmol of 3,5-dichloroaniline, 5.5 mmol of ethyl glyoxalate solution (50%, toluene), 15.0 mmol of 2-fluorostyrene and 5.0 mmol of trifluoroacetic acid in 30.0 ml of acetonitrile. The subsequent saponification was carried out with 1.0 ml of sodium hydroxide solution (6 N, water) in 20.0 ml of ethanol.

- 15 An ESI-MS was run to characterize the product:

MS (EI) m/z: (M\*) 339

**Example 11**

20




5,7-Dichloro-4-(3-fluorophenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid (**11**)

Compound **11** was prepared by the basic process from 5.0 mmol of 3,5-dichloroaniline, 5.5 mmol of ethyl glyoxalate solution (50%, toluene), 15.0 mmol of 3-fluorostyrene and 5.0 mmol of trifluoroacetic acid in 30.0 ml of acetonitrile. The subsequent saponification was carried out with 1.0 ml of sodium hydroxide solution (6 N, water) in 20.0 ml of ethanol.

An ESI-MS was run to characterize the product:

MS (EI) m/z: (M<sup>+</sup>) 340

#### 15 Example 12




5,7-Dichloro-4-(4-fluorophenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid (**12**)

Compound **12** was prepared by the basic process from 5.0 mmol of 3,5-dichloroaniline, 5.5 mmol of ethyl glyoxalate solution (50%, toluene), 15.0 mmol of 4-fluorostyrene and 5.0 mmol of trifluoroacetic acid in 30.0 ml of acetonitrile. The subsequent saponification was carried out with 1.0 ml of sodium hydroxide solution (6 N, water) in 20.0 ml of ethanol.

An ESI-MS was run to characterize the product:

MS (EI) m/z: (M\*) 340

5 **Example 13**




10 5,7-Dichloro-4-(2-chlorophenyl)-1,2,3,4-  
tetrahydroquinoline-2-carboxylic acid (**13**)

Compound **13** was prepared by the basic process from 5.0 mmol  
of 3,5-dichloroaniline, 5.5 mmol of ethyl glyoxalate  
solution (50%, toluene), 15.0 mmol of 2-chlorostyrene and  
15 5.0 mmol of trifluoroacetic acid in 30.0 ml of  
acetonitrile. The subsequent saponification was carried  
out with 1.0 ml of sodium hydroxide solution (6 N, water)  
in 20.0 ml of ethanol.

An ESI-MS was run to characterize the product:

20

MS (EI) m/z: (M\*) 356

**Example 14**

- 5 4-(3-Bromophenyl)-5,7-dichloro-1,2,3,4-tetrahydroquinoline-  
2-carboxylic acid (**14**)


Compound **14** was prepared by the basic process from 5.0 mmol  
of 3,5-dichloroaniline, 5.5 mmol of ethyl glyoxalate  
10 solution (50%, toluene), 15.0 mmol of 3-bromostyrene and  
5.0 mmol of trifluoroacetic acid in 30.0 ml of  
acetonitrile. The subsequent saponification was carried  
out with 1.0 ml of sodium hydroxide solution (6 N, water)  
in 20.0 ml of ethanol.

- 15 An ESI-MS was run to characterize the product:

MS (EI) m/z: (M\*) 401

**Example 15**

20






4-(4-Bromophenyl)-5,7-dichloro-1,2,3,4-tetrahydroquinoline-2-carboxylic acid **(15)**

Compound **15** was prepared by the basic process from 5.0 mmol  
5 of 3,5-dichloroaniline, 5.5 mmol of ethyl glyoxalate  
solution (50%, toluene), 15.0 mmol of 4-bromostyrene and  
5.0 mmol of trifluoroacetic acid in 30.0 ml of  
acetonitrile. The subsequent saponification was carried  
out with 1.0 ml of sodium hydroxide solution (6 N, water)  
10 in 20.0 ml of ethanol.

An ESI-MS was run to characterize the product:


MS (EI) m/z: (M<sup>+</sup>) 401

15 **Example 16**




20 7,8-Dichloro-4-(2-chlorophenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid **(16)**

Compound **16** was prepared by the automated process from 2,3-dichloroaniline, glyoxylic acid and 2-chlorostyrene.

**Example 17**


- 5 6-Cyano-4-(2,3,4-trimethoxyphenyl)-1,2,3,4-  
tetrahydroquinoline-2-carboxylic acid (**17**)

Compound **17** was prepared by the automated process from 4-  
aminobenzonitrile, glyoxylic acid and 2,3,4-  
10 tetramethoxystyrene.

**Example 18**

- 15 6,8,9-Trichloro-2,3,3a,4,5,9b-hexahydrofuro[3,2-c]-  
quinoline-4-carboxylic acid (**18**)


Compound **67** was prepared by the automated process from  
20 2,4,5-trichloroaniline, glyoxylic acid and 2,3-  
dihydrofuran.

**Example 19**

- 5 8-Methoxy-4-(4-methoxyphenyl)-3-methyl-1,2,3,4-  
tetrahydroquinoline-2-carboxylic acid **(19)**

Compound **19** was prepared by the automated process from 2-methoxyaniline, glyoxylic acid and *trans*-anethole.

10


**Example 20**

- 15 5,6,8-Trichloro-4-(4-hydroxyphenyl)-3-methyl-1,2,3,4-  
tetrahydroquinoline-2-carboxylic acid **(20)**

Compound **20** was prepared by the automated process from 2,3,5-trichloroaniline, glyoxylic acid and 2-

propenylphenol.

**Example 21**




5

4-(3,4-Dimethoxyphenyl)-8-iodo-1,2,3,4-tetrahydroquinoline-2-carboxylic acid **(21)**

- 10 Compound **21** was prepared by the automated process from 2-iodoaniline, glyoxylic acid and 3,4-dimethoxystyrene.

**Example 22**




15

6-Iodo-4-(4-methylsulfanylphenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid **(22)**

Compound **22** was prepared by the automated process from 4-iodoaniline, glyoxylic acid and 1-methylsulfanyl-4-vinylbenzene.

5 **Example 23**




10 4-(4-Ethoxy-3-methoxyphenyl)-6-phenoxy-1,2,3,4-tetrahydroquinoline-2-carboxylic acid (**23**)

Compound **23** was prepared by the automated process from 4-phenoxyaniline, glyoxylic acid and 1-ethoxy-2-methoxy-4-vinylbenzene.

15


**Example 24**



4-(2-Ethoxynaphthalen-1-yl)-6-iodo-1,2,3,4-tetrahydroquinoline-2-carboxylic acid (**24**)


Compound **24** was prepared by the automated process from 4-iodoaniline, glyoxylic acid and 2-ethoxy-1-vinylnaphthalene.

**Example 25**



8-Chloro-4-(4-propoxyphenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid (**25**)

Compound **25** was prepared by the automated process from 2-chloroaniline, glyoxylic acid and 4-propoxystyrene.

**Example 26**

- 5 4-(2,4-Dimethoxy-3-methylphenyl)-6-phenoxy-1,2,3,4-  
tetrahydroquinoline-2-carboxylic acid **(26)**

Compound **26** was prepared by the automated process from 4-  
phenoxyaniline, glyoxylic acid and 2,4-dimethoxy-3-  
10 methylstyrene.

Examples 27 to 102 were prepared analogously.

Example	Name
27	4-anthracen-9-yl-6-chloro-8-methyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid
28	6-sec-butyl-4-naphthalen-1-yl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid
29	4-(4-hydroxyphenyl)-3-methyl-8-phenoxy-1,2,3,4-tetrahydroquinoline-2-carboxylic acid
30	8-chloro-6-fluoro-4-naphthalen-2-yl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid
31	4-(4-methoxyphenyl)-3-methyl-6-phenoxy-1,2,3,4-tetrahydroquinoline-2-carboxylic acid
32	6-chloro-8-fluoro-4-m-tolyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid
33	8-chloro-6-fluoro-4-m-tolyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid
34	4-(4-bromophenyl)-6-chloro-8-fluoro-1,2,3,4-tetrahydroquinoline-2-carboxylic acid
35	7,8-dichloro-4-(2,4-dimethylphenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid
36	6-chloro-4-(4-chlorophenyl)-7-trifluoromethyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid
37	4-(2-chlorophenyl)-6-cyano-1,2,3,4-tetrahydroquinoline-2-carboxylic acid
38	6-bromo-8-chloro-4-(2,4-dimethylphenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid
39	6-bromo-4-(2-bromophenyl)-8-chloro-1,2,3,4-tetrahydroquinoline-2-carboxylic acid
40	4-(4-hydroxy-3-methoxyphenyl)-3-methyl-6-methylsulfanyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid
41	6-cyano-3,4-bis(4-methoxyphenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid
42	ethyl 8-chloro-6-fluoro-3,4-bis(4-methoxyphenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylate
43	5,7-dichloro-4-(4-chlorophenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid
44	5,7-dichloro-4-(3-chlorophenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid
45	5,7-dichloro-4-(4-chlorophenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid
46	1,3-dichloro-5,6,6a,7,8,12b-hexahydrobenzo[k]-phenanthridine-6-carboxylic acid
47	1,3-dichloro-5,6a,7,11b-tetrahydro-6H-indeno[2,1-c]-quinoline-6-carboxylic acid
48	5,7-dichloro-4-(3,5-dimethylphenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid
49	7,9-dichloro-2,3,3a,4,5,9b-hexahydro-1H-cyclopenta[c]quinoline-4-carboxylic acid



**Example 50**

Receptor binding (glycine binding site of the NMDA receptor channel)

5

The tests for determining the affinity of the compounds of formula I according to the invention for the glycine binding site of the NMDA receptor channel were carried out on brain membrane homogenates (homogenate of the cortex and hippocampus areas of the brain of male rats, Wistar strain) [B.M. Baron, B.W. Siegel, B.L. Harrison, R.S. Gross, C. Hawes and P. Towers, Journal of Pharmacology and Experimental Therapeutics, vol. 279, p. 62, 1996].

15 For this purpose, the cortex and hippocampus were freed without cutting from freshly removed rat brains, homogenized in 5 mmol/l TRIS acetate buffer, 0.32 mol/l sucrose, pH 7.4 (10 ml/g of fresh weight), using a Potter homogenizer (Braun/Melsungen, 10 piston strokes at 500 rpm), with ice cooling, and then centrifuged for 10 minutes at 1000 g and 4°C. The first supernatant was collected and the sediment was homogenized again with 5 mmol/l TRIS acetate buffer, 0.32 mol/l sucrose, pH 7.4 (5 ml/g of original fresh weight), using the Potter homogenizer (10 piston strokes at 500 rpm), with ice cooling, and centrifuged for 10 minutes at 1000 g and 4°C. The resulting supernatant was combined with the supernatant from the first centrifugation and centrifuged at 17,000 g for 20 minutes at 4°C. The supernatant after this centrifugation was discarded and the membrane sediment was taken up with 5 mmol/l TRIS acetate buffer, pH 8.0 (20 ml/g of original fresh weight), and homogenized with 10 piston strokes at 500 rpm.

35 The membrane homogenate was then incubated for 1 hour at 4°C and centrifuged for 30 minutes at 50,000 g and 4°C. The

supernatant was discarded and the centrifuge tube containing the membrane sediment was sealed with Parafilm and frozen for 24 hours at -20°C. On the following day the membrane sediment was thawed, taken up with ice-cold

5 5 mmol/l TRIS acetate buffer, 0.1% saponin (w/v), pH 7.0 (10 ml/g of original fresh weight), homogenized with 10 piston strokes at 500 rpm and then centrifuged for 20 minutes at 50,000 g and 4°C. The resulting supernatant was discarded and the sediment was taken up in a small volume

10 with 5 mmol/l TRIS acetate buffer, pH 7.0 (approx. 2 ml/g of original fresh weight), and homogenized again with 10 piston strokes at 500 rpm. After determination of the protein content, the membrane homogenate was adjusted to a protein concentration of 10 mg protein/ml with 5 mmol/l

15 TRIS acetate buffer, pH 7.0, and frozen in aliquots until tested.

For the receptor binding test, aliquots were thawed, diluted to 1:10 with 5 mmol/l TRIS acetate buffer, pH 7.0,

20 homogenized with 10 piston strokes at 500 rpm using the Potter homogenizer (10 piston strokes at 500 rpm), with ice cooling, and centrifuged for 60 minutes at 55,000 g and 4°C. The supernatant was decanted and the membrane sediment was adjusted to a protein concentration of 1 mg/ml with ice-

25 cold 50 mmol/l TRIS acetate buffer, pH 7.0, homogenized again with 10 piston strokes at 500 rpm and kept in suspension by stirring on a magnetic stirrer in an ice bath. 100 µl aliquots of this membrane homogenate were used per ml of preparation in the receptor binding test

30 (0.1 mg protein/ml in the final preparation).

In the binding test, 50 mmol/l TRIS acetate buffer, pH 7.0, was used as the buffer and 1 nmol/l (<sup>3</sup>H)-MDL 105.519 (B.M. Baron et al., 1996) was used as the radioactive ligand.

35 The proportion of non-specific binding was determined in the presence of 1 mmol/l glycine.

In other preparations the compounds according to the invention were added in concentration series and the displacement of the radioactive ligand from its specific binding to the glycine binding site of the NMDA receptor channel was measured. The preparations, in triplicate in each case, were incubated for 120 minutes at 4°C and then harvested by means of filtration through glass fibre filter mats (GF/B) in order to determine the radioactive ligand bound to the membrane homogenate. The radioactivity retained on the glass fibre filters was measured in a  $\beta$  counter after the addition of scintillator.

The affinity of the compounds according to the invention for the glycine binding site of the NMDA receptor channel was calculated as the  $IC_{50}$  value (concentration causing 50% displacement of the radioactive ligand from its specific binding) according to the law of mass action by means of non-linear regression and is indicated in Table 1 as the  $K_i$  value (mean of 3 independent experiments) after conversion (according to the Cheng-Prussoff relationship), or as the percentage of the previously bound radioactive ligand (see above) which is displaced from its specific binding by a concentration of 10  $\mu\text{mol/l}$  of the substance according to the invention to be tested.

25

Table 1:

Example	Glycine binding site of the NMDA receptor channel	
	$K_i$ ( $\mu\text{mol/l}$ )	Displacement (%, 10 $\mu\text{mol/l}$ )
2	0.3	100

**Example 51**

NMDA/glycine-induced ionic fluxes on RNA-injected *Xenopus* oocytes

5

The test to determine functional changes of the NMDA receptor channel due to the compound of formula I according to the invention was performed on oocytes of the South African clawed frog, *Xenopus laevis*. This was done by forming neuronal NMDA receptor channels in oocytes after the injection of rat brain RNA and measuring ionic fluxes triggered by the coapplication of NMDA and glycine.

10

Stage V and VI *Xenopus* oocytes (Dumont, J.N., Journal of Morphology, vol. 136, 1972) were microinjected with total RNA from adult rat brain tissue (100-130 ng/cell) and kept at 20°C for up to 10 days in culture medium (composition in mmol/l: 88.0 NaCl, 1.0 KCl, 1.5 CaCl<sub>2</sub>, 0.8 MgSO<sub>4</sub>, 2.4 NaHCO<sub>3</sub>, 5 HEPES, 100 IU/ml penicillin, 100 µg/ml streptomycin, pH 7.4). Transmembrane ionic fluxes were recorded by the conventional two-electrode voltage clamp technique at a holding potential of -70 mV (P. Bloms-Funke, P.M. Madeja, U. Musshoff, E.-J. Speckmann, Neuroscience Letters, vol. 205, p. 115, 1996). The data were plotted and the experimental apparatus controlled using the OTC interface and Cellworks software (npi, FRG). The compounds according to the invention were added to a nominally Mg<sup>2+</sup>-free medium (composition in mmol/l: 89.0 NaCl, 1.0 KCl, 1.8 CaCl<sub>2</sub>, 2.4 NaHCO<sub>3</sub>, 5 HEPES, pH 7.4) and applied systemically by means of a concentration clamp (npi, FRG). To test substance effects mediated via the glycine B binding site of the NMDA receptor channel, the glycine dose-effect curve was plotted with and without the particular compound according to the invention. To do this, NMDA in a fixed concentration of 100 µmol/l was cumulatively coapplied with glycine in increasing

20

25

30

35

concentrations (0-100  $\mu\text{mol/l}$ ). The experiment was then repeated in the same manner with a fixed concentration of the compound according to the invention. To estimate the selectivity for NMDA versus AMPA receptor channels, the effects of the compound according to the invention (10  $\mu\text{mol/l}$ ) was additionally studied on ionic fluxes triggered by AMPA (100  $\mu\text{mol/l}$ ). The current amplitudes were normalized to that of the control response to the coapplication of NMDA (100  $\mu\text{mol/l}$ ) with glycine (10  $\mu\text{mol/l}$ ). The data were analyzed with Igor-Pro software (Version 3.1, WaveMetrics, USA). All the results were expressed as the mean  $\pm$  standard error (SEM) of at least 3 experiments on different oocytes from at least two frogs. The significance for unpaired and paired measurable variables is determined by the Mann-Whitney U-test and the Wilcoxon test (Sysstat, SPSS Inc., USA) respectively.  $\text{EC}_{50}$  values are calculated according to the following formula:

$$Y = Y_{\min} + (Y_{\max} - Y_{\min}) / (1 + (X/\text{EC}_{50})^{-p})$$

( $Y_{\min}$  = minimum test value,  $Y_{\max}$  = maximum test value,  $Y$  = relative current amplitude,  $X$  = concentration of test substance,  $p$  = slope factor). On right shifting of the glycine dose-effect curve, the  $pA_2$  value of the compound according to the invention was determined graphically by means of Schild regression. Concentration ratios were calculated using the  $\text{EC}_{50}$  values, which were determined independently for each dose-effect curve.

The right shifting of the glycine dose-effect curve is shown for Example no. 1 (relative amplitude: current amplitude, normalized to the response after application of NMDA/glycine (100/10  $\mu\text{mol/l}$ )). The results for selected compounds according to the invention in respect of their effects on the glycine dose-effect curve and on AMPA-induced ionic fluxes have been collated in Table 2.

Table 2: Effects of the compounds according to the invention on ionic fluxes triggered by NMDA/glycine and AMPA on RNA-injected oocytes

Example no.	NMDA/glycine-induced ionic fluxes pA <sub>2</sub> value relating to the glycine dose-effect curve	AMPA-induced ionic fluxes Inhibition at 10 µmol/l of the compounds according to the invention
1	6.40	5.4% (n = 2)

5

### Example 52

#### Formalin test on the rat

- 10 The experiments for determining the antinociceptive action of the compounds of formula I according to the invention were performed in the formalin test on male rats (Sprague-Dawley, 150 - 170 g).
- 15 In the formalin test a distinction is made between the first (early) phase (0 - 15 min after the formalin injection) and the second (late) phase (15 - 60 min after the formalin injection) (D. Dubuisson, S.G. Dennis, Pain, 4, 161 - 174 (1977)). The early phase represents a model
- 20 for acute pain as a direct reaction to the formalin injection, while the late phase is regarded as a model for persistent (chronic) pain (T.J. Coderre, J. Katz, A.L. Vaccarino, R. Melzack, Pain, vol. 52, p. 259, 1993).
- 25 The compounds according to the invention were studied in the second phase of the formalin test in order to make predictions about the actions of substances on chronic/inflammatory pain.
- 30 A single subcutaneous formalin injection (50 µl, 5%) in the dorsal side of the right back paw of free-moving

experimental animals induced a nociceptive reaction which is represented by the following behavioural parameters: lifting and holding of the affected paw (score 1), shaking or twitching (score 2), licking and biting (score 3). The differing modes of behaviour triggered by the formalin injection were recorded continuously by observing the animals in the late phase of the formalin test, and differently weighted in an evaluation. Normal behaviour, where the animal distributes its weight evenly over all four paws, was recorded as a score of 0. The time of administration before the formalin injection was chosen according to the mode of administration of the compounds according to the invention (intraperitoneal: 15 min, intravenous: 5 min). After the injection of substances that have an antinociceptive action in the formalin test, the described modes of behaviour (score 1 - 3) of the animals are reduced or even eliminated. The comparison was made with control animals which had received vehicle (solvent) before the administration of formalin. The nociceptive behaviour was calculated as the so-called pain rate (PR). The different behavioural parameters were differently weighted (factor 0, 1, 2, 3). The calculation was performed in subintervals of 3 min according to the following formula:

$$PR = [(T_0 \times 0) + (T_1 \times 1) + (T_2 \times 2) + (T_3 \times 3)]/180$$

$T_0$ ,  $T_1$ ,  $T_2$  and  $T_3$  corresponding to the time in seconds in which the animal exhibited the mode of behaviour 0, 1, 2 or 3, respectively. The number of animals in the substance and vehicle groups,  $n$ , is 10 in each case. Based on the PR calculations, the substance effect was determined as a percentage change relative to the control. The  $ED_{50}$  calculations were performed by regression analysis.

All the compounds according to the invention which were

tested showed a moderately strong to strong inhibition of the formalin-induced nociception.

The results of selected studies in the formalin test on the  
5 rat are collated in the Table below.

Table 3:

Compound	Mode of administration	Dose [mg/kg]	% inhibition of the formalin-induced nociception
1	i.p.	21.5	64.5

10 **Example 53: General preparation of the salts according to the invention from the compounds of one of Examples 1-49**


One equivalent of the compound of one of Examples 1 to 49,  
15 preferably an imino acid or carboxylic acid, is suspended in a small volume of water and one equivalent of 1 N aqueous alkaline solution, preferably NaOH or KOH, is added. If the solubility is poor, methanol is added dropwise until the compound has completely dissolved.  
20 After stirring for 30 minutes at room temperature, the solution is concentrated on a rotary evaporator and the residual solution is frozen at -60°C in an isopropanol/dry ice mixture and freeze-dried. The salts, especially of imino acids and preferably the sodium or potassium salts,  
25 are usually obtained as colourless solids.

An alternative possibility is to prepare the potassium or sodium salts with potassium or sodium trimethylsilanolate (E.D. Laganis, B.L. Chenard, Tetrahedron Letters, 25, 5831-  
30 5834 (1984)). Potassium or sodium trimethylsilanolate is dissolved under nitrogen in an organic solvent (dichloromethane, toluene, THF) and the ester or acid is added all at once. The reaction mixture is stirred for



four hours at room temperature and filtered. The usually colourless solid is washed with diethyl ether and dried under vacuum. The potassium or sodium salts are obtained as solids.


5

**Example 54**

10 7,9-Dichloro-2,3,3a,4,5,9b-hexahydro-1H-cyclopenta[c]-  
quinoline-4-carboxylate; sodium salt **(54)**

7,9-Dichloro-2,3,3a,4,5,9b-hexahydro-1H-cyclopenta[c]-  
quinoline-4-carboxylic acid **(49)** is treated according to

15 Example 106 to give 7,9-dichloro-2,3,3a,4,5,9b-hexahydro-  
1H-cyclopenta[c]quinoline-4-carboxylate; sodium salt **(54)**.

**Example 55**

20

7,9-Dichloro-3a,4,5,9b-tetrahydro-3H-cyclopenta[c]-  
quinoline-4-carboxylate; sodium salt **(55)**

Compound 1 prepared according to Example 1 is treated  
5 according to Example 53 to give 7,9-dichloro-3a,4,5,9b-  
tetrahydro-3H-cyclopenta[c]quinoline-4-carboxylate; sodium  
salt **(55)**.

Molecular weight calculated by ESI-MS: 284.14 g/mol;  
10 measured molecular weight: 282.3 (M-H), 238.4 (M-CO<sub>2</sub>).

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO/TMS<sub>ext</sub>): d = 2.15 - 2.40 ppm (m, 2H, CH<sub>2</sub>);  
3.35 ppm (q, 1H, CH); 3.50 ppm (m, 1H, CH); 4.05 ppm (dd,  
1H, CH); 5.60 ppm (m, 1H, olefin H); 5.70 ppm (m, 1H, NH);  
15 5.80 ppm (M, 1H, olefin CH); 6.60 ppm (d, 1H, aryl CH);  
6.85 ppm (d, 1H, aryl CH).

**Example 56: Receptor binding of the salts according to the  
invention at the glycine binding site of the  
20 NMDA receptor channel**

Compounds **54** and **55** are studied for receptor binding as  
explained in Example 50:

25 Table 4:

Example	Glycine binding site of the NMDA receptor channel	
	Ki (μmol/l)	Displacement (%, 10 μmol/l)
<b>54</b>		<b>97</b>
<b>55</b>	<b>0.35</b>	<b>90</b>

**Example 57: Formalin test**

30 Compound **55** was studied in the formalin test as described  
in Example 52.

Table 5:

Compound	Mode of administration	Dose [mg/kg]	% inhibition of the formalin-induced nociception
55	i.v.	68.1	56

**Example 58: Parenteral mode of administration**

5


38.5 g of compound **55** are dissolved in 1 l of water for injection at room temperature and then adjusted to isotonic conditions by the addition of anhydrous glucose for injection.

10

**Claims**

1. Substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivatives of general formula I:

5



I,

in the form of their physiologically acceptable salts with cations or bases, optionally in the form of their racemates or their pure stereoisomers, especially enantiomers or diastereoisomers, or in the form of mixtures of the stereoisomers, especially the enantiomers or diastereoisomers, in any desired mixing ratio,

15 in which





**either**

20 **R<sup>1</sup>** and **R<sup>2</sup>** together form the following, each of which is monosubstituted or polysubstituted or unsubstituted:

- (CH<sub>2</sub>)<sub>n</sub>-, where n = 3-10,
- CH=CH-CH<sub>2</sub>-, -CH<sub>2</sub>-CH=CH-,
- CH=CH-CH<sub>2</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-CH<sub>2</sub>-CH=CH-,
- 25 -CH<sub>2</sub>-CH=CH-CH<sub>2</sub>-,
- CH<sub>2</sub>-CH=CH-CH<sub>2</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-CH<sub>2</sub>-CH=CH-CH<sub>2</sub>-,
- CH<sub>2</sub>-CH<sub>2</sub>-CH=CH-CH<sub>2</sub>-CH<sub>2</sub>-,
- O-CH<sub>2</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-CH<sub>2</sub>-O-,
- O-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-O-,

-CH<sub>2</sub>-O-CH<sub>2</sub>-,

-CH<sub>2</sub>-CH<sub>2</sub>-O-CH<sub>2</sub>-, -CH<sub>2</sub>-O-CH<sub>2</sub>-CH<sub>2</sub>-,



X = O, S

5

**R<sup>3</sup>** is selected from

10 H; C<sub>1</sub>-C<sub>18</sub>-alkyl, C<sub>2</sub>-C<sub>18</sub>-alkenyl or C<sub>2</sub>-C<sub>18</sub>-alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted; C<sub>3</sub>-C<sub>8</sub>-cycloalkyl  
 15 which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted, or a corresponding heterocycle in which at least one ring C atom is replaced by N, S or O; alkylaryl or alkylheteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted; and aryl or

heteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted;

$R^4$  is selected from

5

$R^{4a}$  or  $ZR^{4a}$ , where  $Z = C_1-C_6$ -alkyl,  $C_2-C_6$ -alkenyl or  $C_2-C_6$ -alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted; and  $R^{4a}$  is selected from

10

H;  $C_1-C_{12}$ -alkyl,  $C_2-C_{12}$ -alkenyl or  $C_2-C_{12}$ -alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted;  $C_3-C_8$ -cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted, or a corresponding heterocycle in which at least one ring C atom is replaced by S, O or N; and aryl or heteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted;

15

20

$C(O)R^9$ ,  $C(O)OR^9$ ,  $C(S)R^9$ ,  $C(S)OR^9$  or  $S(O_2)R^9$ , where  $R^9$  is selected from

25

H;  $C_1-C_{10}$ -alkyl,  $C_2-C_{10}$ -alkenyl or  $C_2-C_{10}$ -alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted;  $C_3-C_8$ -cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted, or a corresponding heterocycle in which at least one ring C atom is replaced by S, O or N; alkylaryl or alkylheteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted; and aryl or heteroaryl, each

30

35

of which is monosubstituted or  
polysubstituted or unsubstituted, especially  
phenethyl, 1-adamantyl, 2-adamantyl, 1-  
naphthyl or 2-naphthyl, 2-, 3- or 4-pyridyl  
or thiazolyl;

5

SR<sup>10</sup>, where R<sup>10</sup> is selected from

aryl or heteroaryl, each of which is  
monosubstituted or polysubstituted or  
unsubstituted;

10

C(O)NR<sup>11</sup>R<sup>12</sup>, C(O)NR<sup>11</sup>NR<sup>12</sup>R<sup>13</sup>, C(NR<sup>11</sup>)NR<sup>12</sup>R<sup>13</sup>,  
C(S)NR<sup>11</sup>R<sup>12</sup> or C(S)NR<sup>11</sup>NR<sup>12</sup>R<sup>13</sup>, where R<sup>11</sup>, R<sup>12</sup> and R<sup>13</sup>  
independently of one another are selected from

15

H; C<sub>1</sub>-C<sub>18</sub>-alkyl, C<sub>2</sub>-C<sub>18</sub>-alkenyl or C<sub>2</sub>-C<sub>18</sub>-  
alkynyl, each of which is branched or  
unbranched and monosubstituted or  
polysubstituted or unsubstituted; C<sub>3</sub>-C<sub>8</sub>-  
cycloalkyl which is saturated or unsaturated  
and monosubstituted or polysubstituted or  
unsubstituted, or a corresponding  
heterocycle in which at least one ring C  
atom is replaced by S, O or N; alkylaryl or  
alkylheteroaryl, each of which is  
monosubstituted or polysubstituted or  
unsubstituted; and aryl or heteroaryl, each  
of which is monosubstituted or  
polysubstituted or unsubstituted;

20

25

30

R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> independently of one another are selected  
from

35

H; F; Cl; Br; I; CN; NO<sub>2</sub>; and C<sub>1</sub>-C<sub>10</sub>-alkyl, C<sub>2</sub>-C<sub>10</sub>-  
alkenyl or C<sub>2</sub>-C<sub>10</sub>-alkynyl, each of which is branched or

unbranched and monosubstituted or polysubstituted or unsubstituted;

5       $OR^{14}$ ,  $OC(O)R^{14}$ ,  $OC(S)R^{14}$ ,  $C(O)R^{14}$ ,  $C(O)OR^{14}$ ,  $C(S)R^{14}$ ,  
     $C(S)OR^{14}$ ,  $SR^{14}$ ,  $S(O)R^{14}$  or  $S(O_2)R^{14}$ , where  $R^{14}$  is selected  
    from

10      H;  $C_1$ - $C_{10}$ -alkyl,  $C_2$ - $C_{10}$ -alkenyl or  $C_2$ - $C_{10}$ -alkynyl,  
    each of which is branched or unbranched and  
    monosubstituted or polysubstituted or  
    unsubstituted;  $C_3$ - $C_8$ -cycloalkyl which is saturated  
    or unsaturated and monosubstituted or  
    polysubstituted or unsubstituted, or a  
15      corresponding heterocycle in which at least one  
    ring C atom is replaced by S, O or N; alkylaryl  
    or alkylheteroaryl, each of which is  
    monosubstituted or polysubstituted or  
    unsubstituted; and aryl or heteroaryl, each of  
20      which is monosubstituted or polysubstituted or  
    unsubstituted;

25       $NR^{15}R^{16}$ ,  $NR^{15}C(O)R^{16}$ ,  $C(NR^{15})NR^{16}R^{17}$ ,  $NR^{15}C(S)R^{16}$ ,  
     $C(S)NR^{15}R^{16}$ ,  $C(S)NR^{15}NR^{16}R^{17}$  or  $S(O_2)NR^{15}R^{16}$ , where  $R^{15}$ ,  $R^{16}$   
    and  $R^{17}$  independently of one another are selected from

30      H; O;  $C_1$ - $C_{18}$ -alkyl,  $C_2$ - $C_{18}$ -alkenyl or  $C_2$ - $C_{18}$ -  
    alkynyl, each of which is branched or unbranched  
    and monosubstituted or polysubstituted or  
    unsubstituted;  $C_3$ - $C_8$ -cycloalkyl which is saturated  
    or unsaturated and monosubstituted or  
    polysubstituted or unsubstituted, or a  
35      corresponding heterocycle in which at least one  
    ring C atom is replaced by S, O or N; alkylaryl  
    or alkylheteroaryl, each of which is  
    monosubstituted or polysubstituted or  
    unsubstituted; and aryl or heteroaryl, each of



which is monosubstituted or polysubstituted or unsubstituted;

or

5

$R^{15}$  and  $R^{16}$  or  $R^{16}$  and  $R^{17}$  together form a  $C_3$ - $C_8$ -cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted, or a corresponding heterocycle in which at least one ring C atom is replaced by S, O or N; or

10

$R^5$  and  $R^6$ ,  $R^6$  and  $R^7$  or  $R^7$  and  $R^8$  together form

15

$=CR^{18}-CH=CH-CH=$  or  $=CH-CR^{18}=CH-CH=$ , where  $R^{18}$  is selected from


H; F; Cl; Br; I; OH; and  $C_1$ - $C_{10}$ -alkyl,  $C_2$ - $C_{10}$ -alkenyl or  $C_2$ - $C_{10}$ -alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted,

20

with the proviso that

25

if  $R^1$  and  $R^2$  together form  $-CH=CH-CH_2-$  or



30


and  $R^3$  is (-)-p-menthan-3-ol, especially menthol or borneol,  $R^7 \neq Cl$  and  $R^5$ ,  $R^6$  and  $R^8 \neq H$  simultaneously,

if  $R^1$  and  $R^2$  together form  $-CH=CH-CH_2-$  and  $R^3$  is  $CH_3$ ,  $R^7 \neq H$ , Cl or  $OCH_3$  and  $R^5$ ,  $R^6$  and  $R^8 \neq H$  simultaneously,

if  $R^{1b}$  and  $R^{2a}$  together form  $-\text{CH}=\text{CH}-\text{CH}_2-$  and  $R^3$  is H,  $R^7 \neq \text{OCH}_3$  or  $\text{C}(\text{O})\text{NH}_2$  and  $R^5$ ,  $R^6$  and  $R^8 \neq \text{H}$ ,  $R^5$  and  $R^7 \neq \text{CH}_3$  and  $R^6$  and  $R^8 \neq \text{H}$ , or  $R^5 \neq \text{OCH}_3$  and  $R^6$ ,  $R^7$  and  $R^8 \neq \text{H}$  simultaneously, or

5

if  $R^{1b}$  and  $R^{2a}$  together form



10 or  $-\text{O}-\text{CH}_2-\text{CH}_2-$  and  $R^3$  is  $\text{C}_2\text{H}_5$ ,  $R^7 \neq \text{H}$ , Cl,  $\text{CH}_3$ ,  $\text{OCH}_3$  or  $\text{NO}_2$  and  $R^5$ ,  $R^6$  and  $R^8 \neq \text{H}$ , or  $R^5 \neq \text{NO}_2$  and  $R^6$ ,  $R^7$  and  $R^8 \neq \text{H}$  simultaneously;

or

15

$R^1$  is selected from

$\text{C}_1$ - $\text{C}_{10}$ -alkyl which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted;  
20  $\text{C}_3$ - $\text{C}_8$ -cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted, or a corresponding heterocycle in which at least one ring C atom is replaced by S, O or N; alkylaryl which is monosubstituted or polysubstituted or  
25 unsubstituted; and aryl which is monosubstituted or polysubstituted or unsubstituted;

$\text{OR}^{19}$ ,  $\text{SR}^{19}$  or  $\text{SO}_2\text{R}^{19}$ , where  $R^{19}$  is selected from

30  $\text{C}_1$ - $\text{C}_{10}$ -alkyl,  $\text{C}_2$ - $\text{C}_{10}$ -alkenyl or  $\text{C}_2$ - $\text{C}_{10}$ -alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted;  $\text{C}_3$ - $\text{C}_8$ -cycloalkyl which is saturated or unsaturated and monosubstituted or

polysubstituted or unsubstituted, or a corresponding heterocycle in which at least one ring C atom is replaced by S, O or N; and alkylaryl, aryl, alkylheteroaryl or heteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted;

$R^2$  is selected from

H;  $C_1$ - $C_{10}$ -alkyl which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted; and phenyl which is monosubstituted or polysubstituted or unsubstituted, where if  $R^2$  is phenyl,  $R^1$  must be aryl, O-aryl or S-aryl;

$R^3$  is selected from

H;  $C_1$ - $C_{18}$ -alkyl,  $C_2$ - $C_{18}$ -alkenyl or  $C_2$ - $C_{18}$ -alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted;  $C_3$ - $C_8$ -cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted, or a corresponding heterocycle in which at least one ring C atom is replaced by N, S or O; alkylaryl or alkylheteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted; and aryl or heteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted;

$R^4$  is selected from

$R^{4a}$  or  $ZR^{4a}$ , where  $Z = C_1$ - $C_6$ -alkyl,  $C_2$ - $C_6$ -alkenyl or  $C_2$ - $C_6$ -alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted; and  $R^{4a}$  is selected from

5 H; C<sub>1</sub>-C<sub>12</sub>-alkyl, C<sub>2</sub>-C<sub>12</sub>-alkenyl or C<sub>2</sub>-C<sub>12</sub>-alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted; C<sub>3</sub>-C<sub>8</sub>-cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted, or a corresponding heterocycle in which at least one ring C atom is replaced by S, O or N; and aryl or heteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted;

10 C(O)R<sup>9</sup>, C(O)OR<sup>9</sup>, C(S)R<sup>9</sup>, C(S)OR<sup>9</sup> or S(O<sub>2</sub>)R<sup>9</sup>, where R<sup>9</sup> is selected from

15 H; C<sub>1</sub>-C<sub>10</sub>-alkyl, C<sub>2</sub>-C<sub>10</sub>-alkenyl or C<sub>2</sub>-C<sub>10</sub>-alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted; C<sub>3</sub>-C<sub>8</sub>-cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted, or a corresponding heterocycle in which at least one ring C atom is replaced by S, O or N; alkylaryl or alkylheteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted; and aryl or heteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted, especially phenethyl, 1-adamantyl, 2-adamantyl, 1-naphthyl or 2-naphthyl, 2-, 3- or 4-pyridyl or thiazolyl;

SR<sup>10</sup>, where R<sup>10</sup> is selected from

35 aryl or heteroaryl, each of which is monosubstituted or polysubstituted or

unsubstituted;

$C(O)NR^{11}R^{12}$ ,  $C(O)NR^{11}NR^{12}R^{13}$ ,  $C(NR^{11})NR^{12}R^{13}$ ,  
5  $C(S)NR^{11}R^{12}$  or  $C(S)NR^{11}NR^{12}R^{13}$ , where  $R^{11}$ ,  $R^{12}$  and  $R^{13}$   
independently of one another are selected from

10 H;  $C_1$ - $C_{18}$ -alkyl,  $C_2$ - $C_{18}$ -alkenyl or  $C_2$ - $C_{18}$ -  
alkynyl, each of which is branched or  
unbranched and monosubstituted or  
polysubstituted or unsubstituted;  $C_3$ - $C_8$ -  
15 cycloalkyl which is saturated or unsaturated  
and monosubstituted or polysubstituted or  
unsubstituted, or a corresponding  
heterocycle in which at least one ring C  
atom is replaced by S, O or N; alkylaryl or  
20 alkylheteroaryl, each of which is  
monosubstituted or polysubstituted or  
unsubstituted; and aryl or heteroaryl, each  
of which is monosubstituted or  
polysubstituted or unsubstituted;

$R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  independently of one another are selected  
from

25 H; F; Cl; Br; I; CN;  $NO_2$ ; and  $C_1$ - $C_{10}$ -alkyl,  $C_2$ - $C_{10}$ -  
alkenyl or  $C_2$ - $C_{10}$ -alkynyl, each of which is branched or  
unbranched and monosubstituted or polysubstituted or  
unsubstituted;

30  $OR^{14}$ ,  $OC(O)R^{14}$ ,  $OC(S)R^{14}$ ,  $C(O)R^{14}$ ,  $C(O)OR^{14}$ ,  $C(S)R^{14}$ ,  
 $C(S)OR^{14}$ ,  $SR^{14}$ ,  $S(O)R^{14}$  or  $S(O_2)R^{14}$ , where  $R^{14}$  is selected  
from

35 H;  $C_1$ - $C_{10}$ -alkyl,  $C_2$ - $C_{10}$ -alkenyl or  $C_2$ - $C_{10}$ -alkynyl,  
each of which is branched or unbranched and  
monosubstituted or polysubstituted or

unsubstituted; C<sub>3</sub>-C<sub>8</sub>-cycloalkyl which is saturated  
or unsaturated and monosubstituted or  
polysubstituted or unsubstituted, or a  
corresponding heterocycle in which at least one  
5 ring C atom is replaced by S, O or N; alkylaryl  
or alkylheteroaryl, each of which is  
monosubstituted or polysubstituted or  
unsubstituted; and aryl or heteroaryl, each of  
which is monosubstituted or polysubstituted or  
10 unsubstituted;

NR<sup>15</sup>R<sup>16</sup>, NR<sup>15</sup>C(O)R<sup>16</sup>, C(NR<sup>15</sup>)NR<sup>16</sup>R<sup>17</sup>, NR<sup>15</sup>C(S)R<sup>16</sup>,  
C(S)NR<sup>15</sup>R<sup>16</sup>, C(S)NR<sup>15</sup>NR<sup>16</sup>R<sup>17</sup> or S(O<sub>2</sub>)NR<sup>15</sup>R<sup>16</sup>, where R<sup>15</sup>, R<sup>16</sup>  
and R<sup>17</sup> independently of one another are selected from  
15

H; O; C<sub>1</sub>-C<sub>18</sub>-alkyl, C<sub>2</sub>-C<sub>18</sub>-alkenyl or C<sub>2</sub>-C<sub>18</sub>-  
alkynyl, each of which is branched or unbranched  
and monosubstituted or polysubstituted or  
unsubstituted; C<sub>3</sub>-C<sub>8</sub>-cycloalkyl which is saturated  
or unsaturated and monosubstituted or  
20 polysubstituted or unsubstituted, or a  
corresponding heterocycle in which at least one  
ring C atom is replaced by S, O or N; alkylaryl  
or alkylheteroaryl, each of which is  
monosubstituted or polysubstituted or  
25 unsubstituted; and aryl or heteroaryl, each of  
which is monosubstituted or polysubstituted or  
unsubstituted;

30 or

R<sup>15</sup> and R<sup>16</sup> or R<sup>16</sup> and R<sup>17</sup> together form a C<sub>3</sub>-C<sub>8</sub>-  
cycloalkyl which is saturated or unsaturated and  
monosubstituted or polysubstituted or  
unsubstituted, or a corresponding heterocycle in  
35 which at least one C atom is replaced by S, O or

N; or

**R**<sup>5</sup> and **R**<sup>6</sup>, **R**<sup>6</sup> and **R**<sup>7</sup> or **R**<sup>7</sup> and **R**<sup>8</sup> together form

5       =CR<sup>18</sup>-CH=CH-CH= or =CH-CR<sup>18</sup>=CH-CH=, where R<sup>18</sup> is  
selected from

10               H; F; Cl; Br; I; OH; and C<sub>1</sub>-C<sub>10</sub>-alkyl, C<sub>2</sub>-C<sub>10</sub>-  
alkenyl or C<sub>2</sub>-C<sub>10</sub>-alkynyl, each of which is  
branched or unbranched and monosubstituted or  
polysubstituted or unsubstituted,

with the proviso that

15   if R<sup>4</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> = H,  
• R<sup>1</sup> ≠ CH<sub>3</sub>, R<sup>3</sup> ≠ H or CH<sub>3</sub> and R<sup>2</sup> and R<sup>5</sup> ≠ H simultaneously;  
or  
• R<sup>1</sup> ≠ unsubstituted phenyl, R<sup>3</sup> ≠ C<sub>2</sub>H<sub>5</sub> and R<sup>2</sup> and R<sup>5</sup> ≠ H  
simultaneously;

20   if R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup> and R<sup>8</sup> = H,  
• R<sup>1</sup> ≠ S-phenyl, R<sup>2</sup> ≠ H, R<sup>7</sup> ≠ Cl and R<sup>3</sup> ≠ CH<sub>3</sub> simultaneously;  
or  
• R<sup>1</sup> ≠ S-2-pyridinyl, R<sup>2</sup> ≠ CH<sub>3</sub>, R<sup>7</sup> ≠ OCH<sub>3</sub> and R<sup>3</sup> ≠ -CH<sub>3</sub>-CH=CH<sub>2</sub>  
25   simultaneously; or

if R<sup>2</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>7</sup> = H and R<sup>6</sup> and R<sup>8</sup> = Cl,  
• R<sup>1</sup> ≠ dioxalane and R<sup>3</sup> ≠ -CH<sub>2</sub>-CH<sub>2</sub>-OH simultaneously.

30   2.   Salts according to Claim 1, characterized in that R<sup>4</sup> is  
selected from

35               H; C<sub>1</sub>-C<sub>10</sub>-alkyl, C<sub>2</sub>-C<sub>10</sub>-alkenyl or C<sub>2</sub>-C<sub>10</sub>-alkynyl, each  
of which is branched or unbranched and monosubstituted  
or polysubstituted or unsubstituted; and C<sub>3</sub>-C<sub>8</sub>-  
cycloalkyl which is saturated or unsaturated and

monosubstituted or polysubstituted or unsubstituted;  
and  
C(O)R<sup>9</sup>, where R<sup>9</sup> is selected from

5           H; C<sub>1</sub>-C<sub>10</sub>-alkyl, C<sub>2</sub>-C<sub>10</sub>-alkenyl or C<sub>2</sub>-C<sub>10</sub>-alkynyl,  
each of which is branched or unbranched and  
monosubstituted or polysubstituted or  
unsubstituted; C<sub>3</sub>-C<sub>8</sub>-cycloalkyl which is saturated  
or unsaturated and monosubstituted or  
10           polysubstituted or unsubstituted; and aryl or  
heteroaryl, each of which is monosubstituted or  
polysubstituted or unsubstituted, especially  
phenethyl, 1-adamantyl, 2-adamantyl, 1-naphthyl  
or 2-naphthyl, 2-, 3- or 4-pyridyl or thiazolyl.

15

3.   Salts according to Claim 1 or 2, characterized in that  
R<sup>4</sup> is selected from

20           H; C<sub>1</sub>-C<sub>10</sub>-alkyl which is unsubstituted or  
monosubstituted or polysubstituted; and phenyl which  
is unsubstituted or monosubstituted or  
polysubstituted, preferably H, CH<sub>3</sub> or C<sub>2</sub>H<sub>5</sub> and  
especially H.

25   4.   Salts according to one of Claims 1 to 3, characterized  
in that R<sup>3</sup> is selected from

30           H; C<sub>1</sub>-C<sub>10</sub>-alkyl, C<sub>2</sub>-C<sub>10</sub>-alkenyl or C<sub>2</sub>-C<sub>10</sub>-alkynyl, each  
of which is branched or unbranched and monosubstituted  
or polysubstituted or unsubstituted; C<sub>3</sub>-C<sub>8</sub>-cycloalkyl  
which is saturated or unsaturated and monosubstituted  
or polysubstituted or unsubstituted, or a  
corresponding heterocycle in which at least one ring C  
atom is replaced by N or O; alkylaryl which is  
35           monosubstituted or polysubstituted or unsubstituted;  
and aryl or heteroaryl, each of which is



monosubstituted or polysubstituted or unsubstituted.

5. Salts according to one of Claims 1 to 4, characterized in that  $R^3$  is selected from


5

H;  $C_1$ - $C_4$ -alkyl which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted; and phenyl, benzyl or phenethyl which is monosubstituted or polysubstituted or unsubstituted, preferably H,  $CH_3$  or  $C_2H_5$  and especially H.

10

6. Salts according to one of Claims 1 to 5, characterized in that  $R^1$  and  $R^2$  together form

15  $-O-CH_2-CH_2-$ ,  $(-CH_2-)_n$  where  $n = 3-6$ , preferably 3 or 6,  $-CH=CH-CH_2-$ ,  $-CH=CH-CH_2-CH_2-$ ,



20 preferably  $-CH=CH-CH_2-$  or  $-CH=CH-CH_2-CH_2-$  and especially  $-CH=CH-CH_2-$ .

7. Salts according to one of Claims 1 to 5, characterized in that  $R^1$  is selected from

25

phenyl, naphthyl or anthracenyl which is unsubstituted or monosubstituted or polysubstituted; and  $OR^{19}$  or  $SR^{19}$ , where  $R^{19}$  is selected from

5 C<sub>1</sub>-C<sub>6</sub>-alkyl which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted; C<sub>3</sub>-C<sub>8</sub>-cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted; and aryl which is monosubstituted or polysubstituted or unsubstituted;

10 preferably anthracenyl, naphthyl or, in particular, phenyl which is unsubstituted or monosubstituted or polysubstituted by a substituent selected from

15 F; Cl; Br; I; methoxy; ethoxy; propoxy; methyl; ethyl; propyl (n-propyl, i-propyl); butyl (n-butyl, i-butyl, t-butyl); carboxyl; nitro; benzyloxy; phenyl; hydroxyl; phenoxy; trifluoromethyl; dioxolyl and SCH<sub>3</sub>;

20 or OR<sup>19</sup> or SR<sup>19</sup>, where R<sup>19</sup> is selected from

25 C<sub>1</sub>-C<sub>4</sub>-alkyl which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted; C<sub>3</sub>-C<sub>8</sub>-cycloalkyl which is saturated or unsaturated and monosubstituted or polysubstituted or unsubstituted; and aryl which is monosubstituted or polysubstituted or unsubstituted;

30 especially unsubstituted phenyl, naphthyl and anthracenyl, O-hydroxyethyl, ethoxynaphthyl, 4-hydroxy-3-methoxyphenyl, 4-propoxyphenyl, 2,3,4-trimethylphenyl, 2,4,5-trimethoxyphenyl, SCH<sub>3</sub>, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 2-bromophenyl, 3-bromophenyl, 4-bromophenyl, 2,6-dichlorophenyl, 4-carboxyphenyl, 3-nitrophenyl, 2,4,6-trimethylphenyl, 2,5-dimethylphenyl, 3,4-dimethoxyphenyl, 4-

35

benzyloxy-3-methoxyphenyl, 3-methylphenyl, 4-methoxyphenyl, 4-biphenyl, 4-methylphenyl, 4-ethoxyphenyl, 2-methylphenyl, 2,4-dimethylphenyl, 2,6-dimethylphenyl, 4-hydroxy-3-methoxyphenyl, 4-methylhydroxyphenyl, 4-hydroxyphenyl, 4-phenoxyphenyl, 4-nitrophenyl, 4-chloromethylphenyl, 4-tert-butylphenyl, 3,5-bis(trifluoromethyl)phenyl, 4-acetoxyphenyl, 4-cyanophenyl, 2-methoxyphenyl, 2,6-difluorophenyl, 2-trifluoromethylphenyl, 3-trifluoromethylphenyl, 4-trifluoromethylphenyl, 3-methoxyphenyl, 2-, 3- or 4-benzyloxyphenyl, S-phenyl or 6-chlorobenzo[1,3]dioxol-5-yl.

8. Salts according to one of Claims 1 to 5 or 7, characterized in that  $R^2$  is selected from

15 H;  $C_1$ - $C_4$ -alkyl which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted; and phenyl which is monosubstituted or polysubstituted or unsubstituted, preferably H, unsubstituted phenyl, 4-methoxyphenyl or  $CH_3$  and especially H.

20

9. Salts according to one of Claims 1 to 8, characterized in that  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  independently of one another are selected from

25 H; F; Cl; Br; I; CN;  $NO_2$ ; and  $C_1$ - $C_{10}$ -alkyl,  $C_2$ - $C_{10}$ -alkenyl or  $C_2$ - $C_{10}$ -alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted;

30  $OR^{14}$ ,  $C(O)R^{14}$ ,  $C(O)OR^{14}$  or  $SR^{14}$ ,  $R^{14}$  being selected from

H;  $C_1$ - $C_{10}$ -alkyl,  $C_2$ - $C_{10}$ -alkenyl or  $C_2$ - $C_{10}$ -alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted;  $C_3$ - $C_8$ -cycloalkyl which is saturated or unsaturated and monosubstituted or

35

polysubstituted or unsubstituted, or a corresponding heterocycle in which at least one ring C atom is replaced by S, O or N; alkylaryl or alkylheteroaryl, each of which is  
5 monosubstituted or polysubstituted or unsubstituted; and aryl or heteroaryl, each of which is monosubstituted or polysubstituted or unsubstituted; and

10  $\text{NR}^{15}\text{R}^{16}$  or  $\text{NR}^{15}\text{C}(\text{O})\text{R}^{16}$ ,  $\text{R}^{15}$  and  $\text{R}^{16}$  independently of one another being selected from

H; O;  $\text{C}_1\text{-C}_{10}$ -alkyl,  $\text{C}_2\text{-C}_{10}$ -alkenyl or  $\text{C}_2\text{-C}_{10}$ -alkynyl, each of which is branched or unbranched  
15 and monosubstituted or polysubstituted or unsubstituted.

10. Salts according to one of Claims 1 to 9, characterized in that  $\text{R}^5$ ,  $\text{R}^6$ ,  $\text{R}^7$  and  $\text{R}^8$  independently of one another are  
20 selected from

H; F; Cl; Br; I; CN;  $\text{NO}_2$ ; and  $\text{C}_1\text{-C}_6$ -alkyl,  $\text{C}_2\text{-C}_6$ -alkenyl or  $\text{C}_2\text{-C}_6$ -alkynyl, each of which is branched or unbranched and monosubstituted or polysubstituted or  
25 unsubstituted;

$\text{OR}^{14}$ ,  $\text{C}(\text{O})\text{R}^{14}$ ,  $\text{C}(\text{O})\text{OR}^{14}$  or  $\text{SR}^{14}$ ,  $\text{R}^{14}$  being selected from

H;  $\text{C}_1\text{-C}_4$ -alkyl which is branched or unbranched and monosubstituted or polysubstituted or  
30 unsubstituted; and aryl which is monosubstituted or polysubstituted or unsubstituted;

preferably,  $\text{R}^5$ ,  $\text{R}^6$ ,  $\text{R}^7$  and  $\text{R}^8$  independently of one another  
35 are selected from

H; F; Cl; Br; I; CN; and C<sub>1</sub>-C<sub>4</sub>-alkyl which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted;

5 OR<sup>14</sup> or SR<sup>14</sup>, R<sup>14</sup> being selected from C<sub>1</sub>-C<sub>4</sub>-alkyl which is branched or unbranched and monosubstituted or polysubstituted or unsubstituted; and aryl which is monosubstituted or polysubstituted or unsubstituted;

10 in particular, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> independently of one another are selected from

15 H; F; Cl; Br; I; CN; CH<sub>3</sub>; CF<sub>3</sub>; t-butyl; i-butyl; -OCH<sub>3</sub>; -OCF<sub>3</sub>; -SCH<sub>3</sub> and -O-phenyl.

11. Salts according to one of Claims 1 to 10, characterized in that

20 R<sup>5</sup>, R<sup>6</sup> and R<sup>8</sup> are H and R<sup>7</sup> is Cl, or R<sup>5</sup> and R<sup>7</sup> are H and R<sup>6</sup> and R<sup>8</sup> are Cl.

12. Salts of substituted 1,2,3,4-tetrahydroquinoline-2-carboxylic acid derivatives according to one of Claims 1 to 25 11, characterized in that they are salts of the following compounds:

30 7,9-dichloro-3a,4,5,9b-tetrahydro-3H-cyclopenta[c]-quinoline-4-carboxylic acid,

8-chloro-3a,4,5,9b-tetrahydro-3H-cyclopenta[c]quinoline-4-carboxylic acid,

35 6-chloro-7-trifluoromethyl-4-(2,4,6-trimethylphenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

4-(2-hydroxyethoxy)-6-trifluoromethoxy-1,2,3,4-  
tetrahydroquinoline-2-carboxylic acid,  
6-iodo-4-(4-methoxyphenyl)-3-methyl-1,2,3,4-  
tetrahydroquinoline-2-carboxylic acid,

5

5,7-dichloro-4-phenyl-1,2,3,4-tetrahydroquinoline-2-  
carboxylic acid,

5,7-dichloro-4-m-tolyl-1,2,3,4-tetrahydroquinoline-2-  
10 carboxylic acid,

5,7-dichloro-4-p-tolyl-1,2,3,4-tetrahydroquinoline-2-  
carboxylic acid,

15 5,7-dichloro-4-(2,4-dimethylphenyl)-1,2,3,4-  
tetrahydroquinoline-2-carboxylic acid,

5,7-dichloro-4-(2-fluorophenyl)-1,2,3,4-  
tetrahydroquinoline-2-carboxylic acid,

20

5,7-dichloro-4-(3-fluorophenyl)-1,2,3,4-  
tetrahydroquinoline-2-carboxylic acid,

5,7-dichloro-4-(4-fluorophenyl)-1,2,3,4-  
25 tetrahydroquinoline-2-carboxylic acid,

5,7-dichloro-4-(2-chlorophenyl)-1,2,3,4-  
tetrahydroquinoline-2-carboxylic acid,

30 5,7-dichloro-4-(4-chlorophenyl)-1,2,3,4-  
tetrahydroquinoline-2-carboxylic acid,

5,7-dichloro-4-(3-bromophenyl)-1,2,3,4-tetrahydroquinoline-  
2-carboxylic acid,

35

5,7-dichloro-4-(4-bromophenyl)-1,2,3,4-tetrahydroquinoline-

2-carboxylic acid,

7,8-dichloro-4-(2-chlorophenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

5

6-cyano-4-(2,3,4-trimethoxyphenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

10 6,8,9-trichloro-2,3,3a,4,5,9b-hexahydrofuro[3,2-c]-quinoline-4-carboxylic acid,

8-methoxy-4-(4-methoxyphenyl)-3-methyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

15 5,6,8-trichloro-4-(4-hydroxyphenyl)-3-methyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

4-(3,4-dimethoxyphenyl)-8-iodo-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

20

6-iodo-4-(4-methylsulfanylphenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

25 4-(4-ethoxy-3-methoxyphenyl)-6-phenoxy-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

4-(2-ethoxynaphthalen-1-yl)-6-iodo-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

30 8-chloro-4-(4-propoxyphenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

4-(2,4-dimethoxy-3-methylphenyl)-6-phenoxy-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

35

4-anthracen-9-yl-6-chloro-8-methyl-1,2,3,4-

tetrahydroquinoline-2-carboxylic acid,

6-sec-butyl-4-naphthalen-1-yl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

5

4-(4-hydroxyphenyl)-3-methyl-8-phenoxy-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

8-chloro-6-fluoro-4-naphthalen-2-yl-1,2,3,4-

10 tetrahydroquinoline-2-carboxylic acid,

4-(4-methoxyphenyl)-3-methyl-6-phenoxy-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

15 6-chloro-8-fluoro-4-m-tolyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

8-chloro-6-fluoro-4-m-tolyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

20

4-(4-bromophenyl)-6-chloro-8-fluoro-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

7,8-dichloro-4-(2,4-dimethylphenyl)-1,2,3,4-

25 tetrahydroquinoline-2-carboxylic acid,

6-chloro-4-(4-chlorophenyl)-7-trifluoromethyl-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

30 4-(2-chlorophenyl)-6-cyano-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

6-bromo-8-chloro-4-(2,4-dimethylphenyl)-1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

35

6-bromo-4-(2-bromophenyl)-8-chloro-1,2,3,4-



tetrahydroquinoline-2-carboxylic acid,

4-(4-hydroxy-3-methoxyphenyl)-3-methyl-6-methylsulfanyl-  
1,2,3,4-tetrahydroquinoline-2-carboxylic acid,

5

6-cyano-3,4-bis(4-methoxyphenyl)-1,2,3,4-  
tetrahydroquinoline-2-carboxylic acid,

5,7-dichloro-4-(4-chlorophenyl)-1,2,3,4-  
10 tetrahydroquinoline-2-carboxylic acid,

5,7-dichloro-4-(3-chlorophenyl)-1,2,3,4-  
tetrahydroquinoline-2-carboxylic acid,

15 5,7-dichloro-4-(4-chlorophenyl)-1,2,3,4-  
tetrahydroquinoline-2-carboxylic acid,

1,3-dichloro-5,6,6a,7,8,12b-hexahydrobenzo[k]-  
phenanthridine-6-carboxylic acid,

20

1,3-dichloro-5,6a,7,11b-tetrahydro-6H-indeno[2,1-c]-  
quinoline-6-carboxylic acid,

5,7-dichloro-4-(3,5-dimethylphenyl)-1,2,3,4-  
25 tetrahydroquinoline-2-carboxylic acid and


7,9-dichloro-2,3,3a,4,5,9b-hexahydro-1H-cyclopenta[c]-  
quinoline-4-carboxylic acid,

30 preference being given to sodium 7,9-dichloro-  
3a,4,5,9b-tetrahydro-3H-cyclopenta[c]quinoline-4-  
carboxylate or sodium 7,9-dichloro-2,3,3a,4,5,9b-  
hexahydro-1H-cyclopenta[c]quinoline-4-carboxylate,  
especially sodium 7,9-dichloro-3a,4,5,9b-tetrahydro-  
35 3H-cyclopenta[c]quinoline-4-carboxylate.

13. Salts according to one of Claims 1 to 12 in the form of the alkali metal salts, preferably the sodium or potassium salts and especially the sodium salts.

- 5 14. Process for the preparation of salts according to Claim 1 in which  $R^4 = H$ , characterized in that anilines of formula II, in which  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  each independently of one another have one of the meanings indicated in Claim 1 or are provided with a protecting group,

10



- glyoxalic acid esters or optionally glyoxalic acid of formula III and olefins of formula IV, in which  $R^1$ ,  $R^2$  and  $R^3$  each independently of one another have one of the meanings indicated in Claim 1 or are provided with a protecting group, are reacted with trifluoroacetic acid at between 0°C and 100°C, any ester groups existing when this basic process has ended being saponified and/or the product formed in the basic process optionally being brought into contact with a strong base, which may already contain the desired cation, in order to form a salt.
- 15
- 20

15. Process according to Claim 14, characterized in that the reaction time is 0.25 - 12 hours, preferably at most 2 h, the reaction is carried out preferably at a temperature of between 20 and 40°C and particularly preferably at room temperature, and/or the reaction is a one-pot reaction.
- 25

16. Process for the preparation of salts according to Claim 1 in which  $R^4 \neq H$ , characterized in that, when the reaction according to Claim 14 has ended, the reaction product in which  $R^4 = H$  is reacted in a consecutive reaction in known manner so that the hydrogen is substituted by  $R^4$  having the other meanings of  $R^4$  in Claim 1.

17. Process according to one of Claims 14 to 16, characterized in that, in a starting material used in the processes, at least one OH group has been replaced by an  $OSi(Ph)_2tert\text{-butyl}$  group, at least one SH group has been replaced by an S-p-methoxybenzyl group and/or at least one  $NH_2$  group has been replaced by an  $NO_2$  group, and at least one and preferably all of the  $OSi(Ph)_2tert\text{-butyl}$  groups are cleaved with tetrabutylammonium fluoride in tetrahydrofuran, and/or at least one and preferably all of the p-methoxybenzyl groups are cleaved with a metal amide, preferably sodamide, and/or at least one and preferably all of the  $NO_2$  groups are preferably all reduced to  $NH_2$  before the end product is purified.

18. Process according to one of Claims 14 to 17, characterized in that, before the end product is purified, a product of the process having at least one  $C(O)OCH_3$  and/or  $C(S)OCH_3$  group, or a product of the process in which  $R^3 = C_{1-4}\text{-alkyl}$ , especially a product of the process in which  $R^3 = CH_3$  or  $C_2H_5$ , is saponified with KOH solution or NaOH solution in methanol or ethanol at  $0^\circ C - 100^\circ C$ , preferably at  $40^\circ C - 60^\circ C$ .

19. Drug containing, as the active ingredient, at least one salt according to one of Claims 1 to 13, optionally in the form of its racemates or its pure stereoisomers, especially enantiomers or diastereoisomers, or in the form of mixtures of the stereoisomers, especially the

enantiomers or diastereoisomers, in any desired mixing ratio, and optionally containing suitable additives and/or auxiliary substances and/or optionally other active ingredients.

5

20. Use of at least one salt according to one of Claims 1 to 13, optionally in the form of its racemates or its pure stereoisomers, especially enantiomers or diastereoisomers, or in the form of mixtures of the stereoisomers, especially  
10 the enantiomers or diastereoisomers, in any desired mixing ratio, for the preparation of a drug for the treatment of pain, especially neuropathic and/or chronic pain, and/or for the treatment of migraine.

15 21. Use of at least one salt according to one of Claims 1 to 13, optionally in the form of its racemates or its pure stereoisomers, especially enantiomers or diastereoisomers, or in the form of mixtures of the stereoisomers, especially the enantiomers or diastereoisomers, in any desired mixing  
20 ratio, for the preparation of a drug for the treatment of urinary incontinence, pruritus, tinnitus aurium and/or diarrhoea.

25 22. Use of at least one salt according to one of Claims 1 to 13, optionally in the form of its racemates or its pure stereoisomers, especially enantiomers or diastereoisomers, or in the form of mixtures of the stereoisomers, especially the enantiomers or diastereoisomers, in any desired mixing  
30 ratio, for the preparation of a drug for the treatment/ prophylaxis of epilepsy, Parkinson's disease, Huntington's chorea, glaucoma, osteoporosis, ototoxicity, withdrawal symptoms associated with alcohol and/or drug abuse, stroke, cerebral ischaemia, cerebral infarction, cerebral oedema, hypoxia and anoxia, and/or for anxiolysis and/or  
35 anaesthesia.

23. Use of at least one salt according to one of Claims 1 to 13, optionally in the form of its racemates or its pure stereoisomers, especially enantiomers or diastereoisomers, or in the form of mixtures of the stereoisomers, especially the enantiomers or diastereoisomers, in any desired mixing ratio, for the preparation of a drug for the treatment/prophylaxis of schizophrenia, Alzheimer's disease, psychosis due to a raised amino acid level, AIDS-related dementia, encephalomyelitis, Tourette syndrome, perinatal asphyxia, inflammatory and allergic reactions, depression, drug and/or alcohol abuse, gastritis, diabetes, cardiovascular disease, respiratory tract disease, cough and/or mental disease.